

# **Anti-ARHGAP18** Antibody

Catalog Number: A08418-1

#### **About ARHGAP18**

ARHGAP18 is one member of the human RhoGAP family with approximately 80 RhoGAP proteins known to be encoded in the human genome. Rho proteins belong to the Ras superfamily that is composed of over 50 members divided into six families, including Ras, Sar, Rho, Ran, Rab and Arf. Rho GTPases are important regulators of the actin cytoskeleton and consequently influence the shape and migration of cells. ARHGAP18 is linked to Ras, and thus, to EGFR-mediated proliferation, migration and differentiation. ARHGAP18 is precisely contained within chromosome 6q22-24, which has been shown to be linked to schizophrenia, suggesting that ARHGEP18 may play a role in this condition.

#### Overview

Product Name	Anti-ARHGAP18 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ARHGAP18 Antibody (Catalog # A08418-1). Tested in ELISA, WB, ICC, IF applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, IF, ICC, WB
Clonality	Polyclonal Clone: SK7
Formulation	ARHGAP18 Antibody is supplied in PBS containing 0.02% sodium azide.
Storage Instructions	ARHGAP18 antibody can be stored at 4°C for three months and -20°C, stable for up to one year. Avoid repeated freeze-thaw cycles. Antibodies should not be exposed to prolonged high temperatures.
Host	Rabbit
Uniprot ID	Q8N392

### **Technical Details**

Immunogen	ARHGAP18 antibody was raised against a 15 amino acid synthetic peptide near the carboxy terminus of human ARHGAP18. The immunogen is located within the last 50 amino acids of ARHGAP18.
Predicted Reactive Species	Chicken
Cross Reactivity	This STEAP2 antibody does not cross-react with other STEAP proteins.
Isotype	IgG
Form	Liquid
Concentration	1 mg/mL






<b>BOSTER</b>
antibody and ELISA experts

Purification	ARHGAP18 Antibody is affinity chromatography purified via peptide column.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  ARHGAP18 antibody can be used for detection of ARHGAP18 by Western blot at 1 - 2 ug/mL.  Antibody can also be used for immunocytochemistry starting at 2.5 ug/mL. For immunofluorescence start at 20 ug/mL.  Antibody validated: Western Blot in mouse samples; Immunocytochemistry in mouse samples and Immunofluorescence in mouse samples. All other applications and species not yet tested. Optimal dilutions for each application should be determined by the researcher.




### Anti-ARHGAP18 Antibody (A08418-1) Images



Western blot analysis of ARHGAP18 in 3T3 cell lysate with ARHGAP18 antibody at (A) 1 and (B) 2 ug/mL.



Immunocytochemistry of ARHGAP18 in 3T3 cells with ARHGAP18 antibody at 2.5 ug/mL.



Immunofluorescence of ARHGAP18 in 3T3 cells with ARHGAP18 antibody at 20 ug/mL.

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-ARHGAP18 Antibody