

Anti-PARP10 Antibody Picoband™

Catalog Number: A08514-2

About PARP10

Poly [ADP-ribose] polymerase 10 is an enzyme that in humans is encoded by the PARP10 gene. Poly(ADP-ribose) polymerases (PARPs), such as PARP10, regulate gene transcription by altering chromatin organization by adding ADP-ribose to histones. PARPs can also function as transcriptional cofactors.

Overview

Product Name	Anti-PARP10 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-PARP10 Antibody Picoband™ catalog # A08514-2. Tested in ELISA, WB, Flow Cytometry applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal 1B9
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q53GL7

Technical Details

Immunogen	E.coli-derived human PARP10 recombinant protein (Position: E14-E251). Human PARP10 shares 76.5% amino acid (aa) sequence identity with mouse PARP10.
Predicted Reactive Species	Bovine, Canine, Chicken, Primate, Sheep, Xenopus, Zebrafish
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross reactivity with other proteins.
Isotype	lgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.25 - $0.5 \mu g/ml$, Human Flow Cytometry (Fixed), 1 - $3 \mu g/1x10^6$ cells, Human ELISA, 0.1 - $0.5 \mu g/ml$, Human
---------------------	---



Anti-PARP10 Antibody Picoband™ (A08514-2) Images




Figure 1. Western blot analysis of PARP10 using anti-PARP10 antibody (A08514-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,
After electrophoresis, proteins were transferred to a
nitrocellulose membrane at 150 mA for 50-90 minutes.
Blocked the membrane with 5% non-fat milk/TBS for 1.5
hour at RT. The membrane was incubated with rabbit antiPARP10 antigen affinity purified polyclonal antibody (Catalog
A08514-2) at 0.5 ug/mL overnight at 4°C, then washed
with TBS-0.1%Tween 3 times with 5 minutes each and
probed with a goat anti-rabbit IgG-HRP secondary antibody
at a dilution of 1:5000 for 1.5 hour at RT. The signal is
developed using an Enhanced Chemiluminescent detection
(ECL) kit (Catalog # EK1002) with Tanon 5200 system. A
specific band was detected for PARP10 at approximately 140
kDa. The expected band size for PARP10 is at 110 kDa.




Figure 2. Flow Cytometry analysis of MCF-7 cells using anti-PARP10 antibody (A08514-2).

Overlay histogram showing MCF-7 cells stained with A08514-2 (Blue line). To facilitate intracellular staining, cells were fixed with 4% paraformaldehyde and permeabilized with permeabilization buffer. The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-PARP10 Antibody (A08514-2, 1 ug/1x10 6 cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10 6 cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10 6) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-PARP10 Antibody