

Anti-OXER1 Antibody Picoband™

Catalog Number: A10420-1

About OXER1

Oxoeicosanoid receptor 1 (OXER1) also known as G-protein coupled receptor 170 (GPR170) is a protein that in humans is encoded by the OXER1 gene located on human chromosome 2p21; it is the principal receptor for the 5-Hydroxyicosatetraenoic acid family of carboxy fatty acid metabolites derived from arachidonic acid. Enables G protein-coupled receptor activity and icosanoid binding activity. Involved in adenylate cyclase-inhibiting G protein-coupled receptor signaling pathway. Predicted to be located in plasma membrane.

Overview

Product Name	Anti-OXER1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-OXER1 Antibody Picoband™ catalog # A10420-1. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q8TDS5

Technical Details

Immunogen	E.coli-derived human OXER1 recombinant protein (Position: M1-Q422).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this

kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blot, 0.25-0.5 µg/ml, Human

Flow Cytometry, 1-3 µg/1x10⁶ cells, Human

Direct ELISA, 0.1-0.5 µg/ml, Human

Anti-OXER1 Antibody Picoband™ (A10420-1) Images




Figure 1. Western blot analysis of OXER1 using anti-OXER1 antibody (A10420-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human THP-1 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-OXER1 antigen affinity purified polyclonal antibody (Catalog # A10420-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for OXER1 at approximately 43-46 kDa. The expected band size for OXER1 is at 43-46 kDa.




Figure 2. Flow Cytometry analysis of U937 cells using anti-OXER1 antibody (A10420-1).

Overlay histogram showing U937 cells stained with A10420-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-OXER1 Antibody (A10420-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-OXER1 Antibody