

Anti-cPLA2 beta/PLA2G4B Antibody Picoband™

Catalog Number: A11137-1

About PLA2G4B

Phospholipase A2, group IVB (cytosolic), also known as PLA2G4B, is a human gene. This gene encodes a member of the cytosolic phospholipase A2 protein family. Phospholipase A2 enzymes hydrolyze the sn-2 bond of phospholipids, releasing lysophospholipids and fatty acids. This enzyme may be associated with mitochondria and early endosomes. Most tissues also express read-through transcripts from the upstream gene into this gene, some of which may encode fusion proteins combining the N-terminus of the upstream gene including its JmjC domain with the almost complete coding region of this gene, including the C2 and cytoplasmic phospholipase A2 domains.

Overview

Product Name	Anti-cPLA2 beta/PLA2G4B Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-cPLA2 beta/PLA2G4B Antibody Picoband™ catalog # A11137-1. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl and 0.2mg Na2HPO4.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	POC869

Technical Details

Immunogen	E.coli-derived human cPLA2 beta/PLA2G4B recombinant protein (Position: N61-H781).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blot, 0.25-0.5ug/ml, Mouse, Rat

Flow Cytometry, 1-3ug/1x10⁶ cells, Human

Direct ELISA, 0.1-0.5ug/ml, Human

Anti-cPLA2 beta/PLA2G4B Antibody Picoband™ (A11137-1) Images




Figure 1. Western blot analysis of cPLA2 beta/PLA2G4B using anti-cPLA2 beta/PLA2G4B antibody (A11137-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30ug of sample under reducing conditions.

Lane 1: rat brain tissue lysates,

Lane 2: mouse brain tissue lysates.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-cPLA2 beta/PLA2G4B antigen affinity purified polyclonal antibody (Catalog # A11137-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for cPLA2 beta/PLA2G4B at approximately 88KD. The expected band size for cPLA2 beta/PLA2G4B is at 88KD.




Figure 2. Flow Cytometry analysis of HeLa cells using anti-cPLA2 beta/PLA2G4B antibody (A11137-1).

Overlay histogram showing HeLa cells stained with A11137-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-cPLA2 beta/PLA2G4B Antibody (A11137-1, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-cPLA2 beta/PLA2G4B Antibody