

Anti-Nucleolar protein of 40 kDa ZCCHC17 Antibody

Catalog Number: A12008-1

About ZCCHC17

Pore-forming (alpha) subunit of voltage-gated potassium channel. Elicits an outward current with fast inactivation. Channel properties may be modulated by cAMP and subunit assembly.

Miyake A., J. Biol. Chem. 274:25018-25025(1999).

Nagase T., DNA Res. 6:337-345(1999).

Overview

Product Name	Anti-Nucleolar protein of 40 kDa ZCCHC17 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Nucleolar protein of 40 kDa ZCCHC17 Antibody catalog # A12008-1. Tested in WB applications. This antibody reacts with Human,Mouse,Rat.
Application	WB
Clonality	Polyclonal
Formulation	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q9NP64

Technical Details

Immunogen	Synthesized peptide derived from human Pbx 4
Predicted Reactive Species	Chimpanzee, Drosophila, Macaque
Isotype	IgG
Form	Liquid
Concentration	1 mg/ml
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.
Some PubMed article(s) citing the expression level of this target are as follows:
Boster Bio's internal QC testing used:
WB: 1:500-1:1000

Anti-Nucleolar protein of 40 kDa ZCCHC17 Antibody (A12008-1) Images




Figure 1. Western blotting validation for Anti-Nucleolar protein of 40 kDa ZCCHC17 Antibody A12008-1

Western blot (WB) analysis of ZCCHC17 polyclonal antibody at 1:500 dilution Lane1:HEK293T whole cell lysate Lane2:RAW264.7 whole cell lysate Lane3:H9C2 whole cell lysate

Electrophoresis was performed on a SDS-PAGE gel. To determine SDS-PAGE gel concentration

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Nucleolar protein of 40 kDa ZCCHC17 Antibody