

Anti-SNX31 Antibody Picoband™

Catalog Number: A14470-1

About SNX31

Predicted to enable phosphatidylinositol binding activity. Predicted to be involved in intracellular protein transport. Predicted to be located in cytoskeleton. Predicted to be part of protein-containing complex. Predicted to be active in early endosome.

Overview

Product Name	Anti-SNX31 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-SNX31 Antibody Picoband™ catalog # A14470-1. Tested in ELISA, Flow Cytometry, WB applications. This antibody reacts with Human.
Application	ELISA, Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ .
Storage Instructions	At -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freezing and thawing.
Host	Rabbit
Uniprot ID	Q8N9S9

Technical Details

Immunogen	E.coli-derived human SNX31 recombinant protein (Position: M1-Q243).
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the

optimal dilution ratio for your samples.
Some PubMed article(s) citing the expression level of this target are as follows:
Boster Bio's internal QC testing used:
Western blot, 0.25-0.5 µg/ml, Human
Flow Cytometry, 1-3 µg/1x10⁶ cells, Human
Direct ELISA, 0.1-0.5 µg/ml, Human

Anti-SNX31 Antibody Picoband™ (A14470-1) Images




Figure 1. Western blot analysis of SNX31 using anti-SNX31 antibody (A14470-1).
Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.
Lane 1: human RT4 whole cell lysates.
After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SNX31 antigen affinity purified polyclonal antibody (Catalog # A14470-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SNX31 at approximately 51 kDa. The expected band size for SNX31 is at 51 kDa.




Figure 2. Flow Cytometry analysis of U251 cells using anti-SNX31 antibody (A14470-1).
Overlay histogram showing U251 cells stained with A14470-1 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-SNX31 Antibody (A14470-1, 1 ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SNX31 Antibody