

# Anti-MCT14 SLC16A14 Antibody

Catalog Number: A15084

#### **About SLC16A14**

Histones are the main constituents of the protein part of chromosomes of eukaryotic cells. They are rich in the amino acids arginine and lysine and have been greatly conserved during evolution. Histones pack the DNA into tight masses of chromatin. Two core histones of each class H2A, H2B, H3 and H4 assemble and are wrapped by 146 base pairs of DNA to form one octameric nucleosome. Histones play a internal role in the regulation of transcription, DNA repair, DNA replication and chromosomal stability. These different functions are established via a complex set of post-translational modifications which either directly or indirectly alter chromatin structure and DNA accessibility to facilitate transcriptional activation or repression or other nuclear processes. Anti-Histone H4 pan Antibody is ideal for research in Chromatin Remodeling, Gene Expression and Epigenetics.

#### Overview

Product Name	Anti-MCT14 SLC16A14 Antibody
Reactive Species	Human, Mouse
Description	Boster Bio Anti-MCT14 SLC16A14 Antibody catalog # A15084. Tested in WB applications. This antibody reacts with Human, Mouse.
Application	WB
Clonality	Polyclonal
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q7RTX9

#### **Technical Details**

Immunogen	Synthesized peptide derived from the Internal region of human MCT14. at AA rangle: 100-180
Predicted Reactive Species	Bovine, Canine, Equine, Guinea Pig, Rabbit, Zebrafish
Isotype	lgG
Form	Liquid
Concentration	This antibody's concentration is >0.5mg/ml.
Purification	Immunogen affinity purified




# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  WB, 1:500-1:2000
---------------------	--



## Anti-MCT14 SLC16A14 Antibody (A15084) Images



Western Blot (WB) analysis of HepG2 cells using MCT14 Polyclonal antibody.

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-MCT14 SLC16A14 Antibody