

Anti-Survivin birc5.1-a Antibody

Catalog Number: A30032


Overview

Product Name	Anti-Survivin birc5.1-a Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Survivin birc5.1-a Antibody (Catalog# A30032). Tested in IP, IHC, WB application(s). This antibody reacts with Human, Mouse, Rat.
Application	IP, IHC, WB
Clonality	Polyclonal
Formulation	50 ul sera
Storage Instructions	Store the antibody at 4°C, stable for 6 months. For long-term storage, store at -20°C. Avoid repeated freeze and thaw cycles.
Host	Rabbit
Uniprot ID	O15392


Technical Details

Immunogen	A Full-length recombinant protein of human Survivin
Predicted Reactive Species	Mouse, Rat
Isotype	Rabbit IgG
Form	Liquid
Concentration	0.5-1mg/ml, actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>WB: 1:1000-1:2000</p> <p>IHC (paraffin): 1:1000-1:5000</p> <p>IHC (frozen): Users should optimize</p> <p>IP: 1:50-1:200</p>


Anti-Survivin birc5.1-a Antibody (A30032) Images



Immunohistochemical analysis of Survivin in a formalin-fixed, paraffin-embedded human brain tissue microarray using primary antibody dilution ratio of 1:2000. A. Differential expression of Survivin in brain tumors from six different patients. A1-A3, high magnification from A. A1, gemistocytic astrocytoma. A2, medulloblastoma. A3, choroid plexus papilloma. Hematoxylin-eosin counterstain.



Immunohistochemical analysis of Survivin in formalin-fixed, paraffin-embedded human prostate and breast cancer using primary antibody dilution ratio of 1:2000. A and A1, prostate cancer at low and high magnification, respectively. B and B1, breast cancer. B1 is a high magnification of an area of metastasis. Hematoxylin-eosin counterstain.



Western blot analysis of Survivin using primary antibody dilution ratio of 1:2000. Survivin was detected in the two tumor cell lines, Jurkat (T cell lymphoma) and RS11846 (non-Hodgkin's lymphoma), but not in lysates from normal tissues. Survivin RP, full-length recombinant human survivin protein was used as a positive control.

9 Publications Citing This Product

1. PubMed ID: 10.3390/molecules18089382, Reactive Oxygen Species Mediate Isoalantolactone-Induced Apoptosis in Human Prostate Cancer Cells
2. PubMed ID: 10.1007/s00432-008-0391-5, Prognostic significance of Survivin and CD44v6 in laryngeal cancer surgical margins
3. PubMed ID: 10.1016/j.envpol.2018.08.003, Insight into the negative impact of ionic liquid: A cytotoxicity mechanism of 1-methyl-3-octylimidazolium bromide

Visit bosterbio.com/anti-survivin-antibody-a30032-boster.html to see all 9 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Survivin birc5.1-a Antibody