

Anti-p53 TP53 Rabbit Monoclonal Antibody

Catalog Number: M00001-1

About TP53

The ion channels activated by glutamate are typically divided into two classes. Those that are sensitive to N-methyl-D-aspartate (NMDA) are designated NMDA receptors (NMDAR) while those activated by alpha-amino-3-hydroxy-5-methyl-4-isoxalone propionic acid (AMPA) are known as AMPA receptors (AMPA). The AMPAR are comprised of four distinct glutamate receptor subunits designated (GluR1-4) and they play key roles in virtually all excitatory neurotransmission in the brain (Keinänen et al., 1990; Hollmann and Heinemann, 1994). The GluR1 subunit is widely expressed throughout the nervous system. Phosphorylation of Ser-845 on GluR1 is thought to be mediated by PKA and phosphorylation of this site increases the conductance of the AMPAR (Roche et al., 1996; Banke et al., 2000). In addition, phosphorylation of this site has been linked to synaptic plasticity as well as learning and memory (Lee et al., 2003; Esteban et al., 2003).

Overview

Product Name	Anti-p53 TP53 Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-p53 TP53 Rabbit Monoclonal Antibody catalog # M00001-1. Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.
Application	IP, IF, IHC, ICC, WB
Clonality	Monoclonal AGC-20
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P04637

Technical Details

Immunogen	A synthesized peptide derived from human p53
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this

kit.

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:


WB 1:500-1:2000

IHC 1:50-1:200


ICC/IF 1:50-1:200

IP 1:50


Anti-p53 TP53 Rabbit Monoclonal Antibody (M00001-1) Images




All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.




Immunohistochemical analysis of paraffin-embedded human colon cancer, using p53 Antibody.



All lanes use the Antibody at 1:3K dilution for 1 hour at room temperature.



Immunofluorescent analysis of HeLa cells, using p53 Antibody .



Western blot analysis of p53 expression in T47D cell lysate.

1. PubMed ID: -, Jiang X, Yuan J, Dou Y, Zeng D, Xiao S. Lipopolysaccharide Affects the Proliferation and Glucose Metabolism of Cervical Cancer Cells Through the FRA1/MDM2/p53 Pathway. *Int J Med Sci* 2021;18(4): 1030-1038. doi:10.7150/ijms.47360.
2. PubMed ID: 33281970, Wang P, Wang C, Liu C. Antitumor effects of dioscin in A431 cells via adjusting ATM/p53-mediated cell apoptosis, DNA damage and migration. *Oncol Lett*. 2021 Jan;21(1):59. doi:10.3892/ol.2020.12321. Epub 2020 Nov 19. PMID:33281970; PMCID:PMC7709553.
3. PubMed ID: 25395712, Li W, Wu D, Wei B, Wang S, Sun H, Li X, Zhang F, Zhang C, Xin Y. Afr J Tradit Complement Altern Med. 2014 Aug 23;11(5):99-104. Ecollection 2014. Anti-Tumor Effect Of Cactus Polysaccharides On Lung Squamous Carcinoma Cells (Sk-Mes-1).

Visit bosterbio.com/anti-p53-rabbit-monoclonal-antibody-m00001-1-boster.html to see all 11 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-p53 TP53 Rabbit Monoclonal Antibody