

Anti-p27 KIP 1 CDKN1B Rabbit Monoclonal Antibody

Catalog Number: M00173-1

About CDKN1B

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

Overview

Product Name	Anti-p27 KIP 1 CDKN1B Rabbit Monoclonal Antibody
Reactive Species	Human, Rat
Description	Boster Bio Anti-p27 KIP 1 CDKN1B Rabbit Monoclonal Antibody catalog # M00173-1. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Rat.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal CGA-3
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P46527

Technical Details

Immunogen	A synthesized peptide derived from human p27 KIP 1
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used:




BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com


WB 1:500-1:2000
IHC 1:50-1:200
ICC/IF 1:50-1:200
IP 1:50
FC 1:50




Anti-p27 KIP 1 CDKN1B Rabbit Monoclonal Antibody (M00173-1) Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Immunohistochemical analysis of paraffin-embedded human colon, using p27 KIP 1 Antibody.




All lanes use the Antibody at $1:1\mbox{K}$ dilution for 1 hour at room temperature.



Immunofluorescent analysis of Hela cells, using p27 KIP 1 Antibody .

Western blot analysis of p27 KIP 1 expression in MCF-7 cell lysate.





1 Publications Citing This Product

1. PubMed ID: 21575146, Hepatitis B virus induces G1 phase arrest by regulating cell cycle genes in HepG2.2.15 cells

Visit bosterbio.com/anti-p27-kip-1-rabbit-monoclonal-antibody-m00173-1-boster.html to see all 1 publications.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-p27 KIP 1 CDKN1B Rabbit Monoclonal Antibody