

Anti-CDK1/Cdc2 Rabbit Monoclonal Antibody

Catalog Number: M00209-4

About CDK1

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

Overview


Product Name	Anti-CDK1/Cdc2 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CDK1/Cdc2 Rabbit Monoclonal Antibody catalog # M00209-4. Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human, Mouse, Rat.
Application	IP, IF, IHC, ICC, WB
Clonality	Monoclonal CBB-3
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P06493

Technical Details


Immunogen	A synthesized peptide derived from human CDK1
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used:

	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50
--	---


Anti-CDK1/Cdc2 Rabbit Monoclonal Antibody (M00209-4) Images



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Western blot analysis of Phospho-CDK1/2/3 (T14) expression in (1) A431 cell lysate; (2) Hela cell lysate.

5 Publications Citing This Product

1. PubMed ID: 33292254, Wang C, Shao S, Deng L, Wang S, Zhang Y. LncRNA SNHG12 regulates the radiosensitivity of cervical cancer through the miR-148a/CDK1 pathway. Cancer Cell Int. 2020 Dec 1;20(1):554. doi:10.1186/s12935-020-01654-5. PMID:33292254; PMCID:PMC7708190.
2. PubMed ID: 30159255, Coxsackievirus A6 induces cell cycle arrest in G0/G1 phase for viral production
3. PubMed ID: 28229049, Human enterovirus 68 interferes with the host cell cycle to facilitate viral production

Visit bosterbio.com/anti-cdk1-rabbit-monoclonal-antibody-m00209-4-boster.html to see all 5 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CDK1/Cdc2 Rabbit Monoclonal Antibody