

## **Anti-PAK1 Monoclonal Antibody**

Catalog Number: M00454

#### **About PAK1**

C3 plays a central role in the activation of the complement system. Its processing by C3 convertase is the central reaction in both classical and alternative complement pathways. After activation C3b can bind covalently, via its reactive thioester, to cell surface carbohydrates or immune aggregates.

#### Overview

| Product Name         | Anti-PAK1 Monoclonal Antibody   |
|----------------------|---|
| Reactive Species     | Human, Mouse, Rat   |
| Description          | Boster Bio Anti-PAK1 Monoclonal Antibody catalog # M00454. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat. |
| Application          | Flow Cytometry, IP, IF, IHC, ICC, WB  |
| Clonality            | Monoclonal ADFA-16  |
| Formulation          | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.   |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.                          |
| Host                 | Rabbit  |
| Uniprot ID           | Q13153  |

#### **Technical Details**

| Immunogen           | A synthesized peptide derived from human PAK1 The activated kinase acts on a variety of targets. Likely to be the GTPase effector that links the Rho-related GTPases to the JNK MAP kinase pathway. Activated by CDC42 and RAC1. Involved in dissolution of stress fibers and reorganization of focal complexes. Involved in regulation of microtubule biogenesis through phosphorylation of TBCB. |
|---------------------|--|
| Isotype             | Rabbit IgG   |
| Form                | Liquid   |
| Concentration       | Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.   |
| Purification        | Affinity-chromatography  |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  |



# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:


Boster Bio's internal QC testing used:

WB 1:500-1:2000 IHC 1:100-1:500 ICC/IF 1:50-1:200 IP 1:40

FC 1:30



### Anti-PAK1 Monoclonal Antibody (M00454) Images



Western blot analysis of PAK1 expression in A431 cell lysate.

#### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-PAK1 Monoclonal Antibody