

Anti-CASR Antibody Picoband™ (monoclonal, 11E9)

Catalog Number: M00574

About CASR

The calcium-sensing receptor (CaSR) is a G protein-coupled receptor that is expressed in the parathyroid hormone (PTH)-producing chief cells of the parathyroid gland, and the cells lining the kidney tubule. It senses small changes in circulating calcium concentration and couples this information to intracellular signaling pathways that modify PTH secretion or renal cation handling, thus this protein plays an essential role in maintaining mineral ion homeostasis. Mutations in this gene cause familial hypocalciuric hypercalcemia, familial, isolated hypoparathyroidism, and neonatal severe primary hyperparathyroidism.

Overview

Product Name	Anti-CASR Antibody Picoband™ (monoclonal, 11E9)
Reactive Species	Human
Description	Boster Bio Anti-CASR Antibody Picoband™ (monoclonal, 11E9) catalog # M00574. Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human.
Application	Flow Cytometry, IHC, ICC, WB
Clonality	Monoclonal 11E9
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P41180

Technical Details

Immunogen	E. coli-derived human CASR recombinant protein (Position: Q926-S1078). Human CASR shares 80.5% and 78.6% amino acid (aa) sequence identity with mouse and rat CASR, respectively.
Predicted Reactive Species	Hepatitis Virus
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot, and HRP Conjugated anti-Mouse IgG Super Vision Assay Kit (SV0001-1) for IHC(P), IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml Immunohistochemistry (Frozen Section), 0.5-1ug/ml Immunocytochemistry, 0.5-1ug/ml Flow Cytometry, 1-3ug/1x10⁶ cells

Anti-CASR Antibody Picoband™ (monoclonal, 11E9) (M00574) Images




Figure 1. Flow Cytometry analysis of NEURO-2A cells using anti-CASR antibody (M00574).
Overlay histogram showing NEURO-2A cells stained with M00574 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-CASR Antibody (M00574, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight® 488 conjugated goat anti-mouse IgG (BA1126, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was mouse IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.




Figure 2. Western blot analysis of CASR using anti-CASR antibody (M00574).
Electrophoresis was performed on a 8% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.
Lane 1: human Hela whole cell lysate,
Lane 2: human A549 whole cell lysate,
Lane 3: human 22RV1 whole cell lysate,
Lane 4: human HepG2 whole cell lysate,
Lane 5: human Caco-2 whole cell lysate.
After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-CASR antigen affinity purified monoclonal antibody (Catalog # M00574) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-CASR Antibody (monoclonal, 11E9)