

Anti-CD68 (Macrophage Marker) Monoclonal Antibody

Catalog Number: M00602

About CD68

This antibody recognizes a glycoprotein of 110kDa, which is identified as CD68. It is important for identifying macrophages in tissue sections. It stains macrophages in a wide variety of human tissues, including Kupffer cells and macrophages in the red pulp of the spleen, in lamina propria of the gut, in lung alveoli, and in bone marrow. It reacts with myeloid precursors and peripheral blood granulocytes. It also reacts with plasmacytoid T cells, which are supposed to be of monocyte/macrophage origin. It shows strong granular cytoplasmic staining of chronic and acute myeloid leukemia and also reacts with rare cases of true histiocytic neoplasia. Lymphomas are negative or show few granules.

Overview

Product Name	Anti-CD68 (Macrophage Marker) Monoclonal Antibody
Reactive Species	Human, Monkey, Rabbit, Cat
Description	Boster Bio Anti-CD68 (Macrophage Marker) Monoclonal Antibody (Catalog # M00602). Tested in Flow Cytometry, IF, IHC applications. This antibody reacts with Human, Monkey, Rabbit, Cat.
Conjugate	Biotin
Application	Flow Cytometry, IF, IHC
Clonality	Monoclonal Clone: SPM130
Formulation	Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.
Storage Instructions	Antibody with azide - store at 2 to 8°C. Antibody without azide - store at -20 to -80°C. Antibody is stable for 24 months. Non-hazardous. No MSDS required.
Host	Mouse
Uniprot ID	P34810

Technical Details

Immunogen	Subcellular fraction of human alveolar macrophages
Predicted Reactive Species	Bovine, Canine, Mouse, Orangutan, Pig, Rabbit, Rat, Deer
Cross Reactivity	Does not cross-react with primate, avian or amphibian GR.
Isotype	IgG1, kappa
Form	Liquid
Concentration	Purified antibody with BSA and azide at 200ug/ml






888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A/G.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Flow Cytometry (1-2ug/million cells) Immunofluorescence (1-2ug/ml) Immunohistochemistry (Formalin-fixed) (1-2ug/ml for 30 minutes at RT) (Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95°C followed by cooling at RT for 20 minutes) Optimal dilution for a specific application should be determined.



Anti-CD68 (Macrophage Marker) Monoclonal Antibody (M00602) Images



Formalin-fixed, paraffin-embedded human Tonsil stained with Anti-CD68 Mouse Monoclonal Antibody (SPM130).

16 Publications Citing This Product

1. PubMed ID: 10.3389/fphar.2018.00618, UPLC/Q-TOFMS-Based Metabolomics Approach to Reveal the Protective Role of Other Herbs in An-Gong-Niu-Huang Wan Against the Hepatorenal Toxicity of Cinnabar and Realgar

2. PubMed ID: 34020949,

Myeloid-derived growth factor inhibits inflammation and alleviates endothelial injury and atherosclerosis in mice

Authors: Meng B,Li Y,Ding Y,Xu X,Wang L,Guo B,Zhu B,Zhang J,Xiang L,Dong J,Liu M,Xiang L,Xiang G.

3. PubMed ID: 32964311, Paschalidi P,Gkouveris I,Soundia A,Kalfarentzos E,Vardas E, Georgaki M,Kostakis G,Erovic BM,Tetradis S,Perisanidis C,Nikitakis NG.The role of M1 and M2 macrophage polarization in progression of medication-related osteonecrosis of the jaw.Clin Oral Investig.2020 Sep 22:10.1007/s00784-020-03602-z.doi:10.1007/s00784-020-03602-z.Epub ahead of print.PMID:32964311;PMCID:PMC7982347.

Visit bosterbio.com/anti-cd68-macrophage-marker-antibody-m00602-boster.html to see all 16 publications.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-CD68 (Macrophage Marker) Monoclonal Antibody