

Anti-ADAM17/Tace Rabbit Monoclonal Antibody

Catalog Number: M00604

About ADAM17

Dynamin-related GTPase required for mitochondrial fusion and regulation of apoptosis. May form a diffusion barrier for proteins stored in mitochondrial cristae. Proteolytic processing in response to intrinsic apoptotic signals may lead to disassembly of OPA1 oligomers and release of the caspase activator cytochrome C (CYCS) into the mitochondrial intermembrane space.

Overview


Product Name	Anti-ADAM17/Tace Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ADAM17/Tace Rabbit Monoclonal Antibody catalog # M00604. Tested in WB, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IP, IF, ICC, WB
Clonality	Monoclonal AOIO-1
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P78536

Technical Details


Immunogen	A synthesized peptide derived from human ADAM17
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used:

	WB 1:500-1:2000 ICC/IF 1:50-1:200 IP 1:50 FC 1:100
--	---


Anti-ADAM17/Tace Rabbit Monoclonal Antibody (M00604) Images




All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Immunofluorescent analysis of HeLa cells, using ADAM17 Antibody.



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of ADAM17 expression in HeLa cell lysate.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ADAM17/Tace Rabbit Monoclonal Antibody