

## Anti-Collagen I COL1A1 Rabbit Monoclonal Antibody

Catalog Number: M00624

### About COL1A1

Dynamin-related GTPase required for mitochondrial fusion and regulation of apoptosis. May form a diffusion barrier for proteins stored in mitochondrial cristae. Proteolytic processing in response to intrinsic apoptotic signals may lead to disassembly of OPA1 oligomers and release of the caspase activator cytochrome C (CYCS) into the mitochondrial intermembrane space.

### Overview


|                      |   |
|----------------------|---|
| Product Name         | Anti-Collagen I COL1A1 Rabbit Monoclonal Antibody   |
| Reactive Species     | Human   |
| Description          | Boster Bio Anti-Collagen I COL1A1 Rabbit Monoclonal Antibody catalog # M00624. Tested in WB, IHC applications. This antibody reacts with Human. |
| Application          | IHC, WB   |
| Clonality            | Monoclonal AFD-3  |
| Formulation          | Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.                             |
| Storage Instructions | Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.      |
| Host                 | Rabbit  |
| Uniprot ID           | P02452/P08123   |

### Technical Details

|                     |  |
|---------------------|--|
| Immunogen           | A synthesized peptide derived from human Collagen I  |
| Isotype             | Rabbit IgG   |
| Form                | Liquid   |
| Concentration       | Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.   |
| Purification        | Affinity-chromatography  |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.<br>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.<br>Some PubMed article(s) citing the expression level of this target are as follows:<br>Boster Bio's internal QC testing used: |

|  |                                   |
|--|-----------------------------------|
|  | WB 1:500-1:2000<br>IHC 1:50-1:200 |
|--|-----------------------------------|

## Anti-Collagen I COL1A1 Rabbit Monoclonal Antibody (M00624) Images



Western blot analysis of Collagen I in Human stomach tissue lysate.




Figure 2. IHC analysis of Collagen I using anti-Collagen I antibody (M00624). Collagen I was detected in a paraffin-embedded section of human breast cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Collagen I Antibody (M00624) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 3. IHC analysis of Collagen I using anti-Collagen I antibody (M00624). Collagen I was detected in a paraffin-embedded section of human colorectal adenocarcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Collagen I Antibody (M00624) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 4. IHC analysis of Collagen I using anti-Collagen I antibody (M00624). Collagen I was detected in a paraffin-embedded section of human lung squamous cell carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Collagen I Antibody (M00624) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 5. IHC analysis of Collagen I using anti-Collagen I antibody (M00624).

Collagen I was detected in a paraffin-embedded section of human placenta tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-Collagen I Antibody (M00624) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Collagen I COL1A1 Rabbit Monoclonal Antibody