

Anti-LAMP1 Monoclonal Antibody

Catalog Number: M00780-1

About LAMP1

Lysosme associated membrane proteins, or LAMP1 and LAMP2, are major constituents of the lysosomal membrane. The two have closely related structures, with 37% sequence homology (2). They are both transmembrane glycoproteins that are localized primarily in lysosomes and late endosomes. Newly synthesized molecules are mostly transported from the trans-Golgi network directly to endosomes and then to lysosomes. A second pathway involves the lamps being delivered from the Golgi to the cell surface, and then along the endocytic pathway to the lysosomes. A minor pathway involves transport via the plasma membrane (3). Upon stimulation, a rapid translocation of intracellular LAMPs to the cell membrane is dependent on a carboxyl-terminal tyrosine ba based motif (YXXI) (1). If there is a disturbance in this spacing, lysosome localization of LAMP1 is abolished and the mutant protein then cycles between the membrane and the endosome (3). This stimulation has also been shown to have an associated release of histamine, leukotriene C (4) and prostaglandin D (2), which shows that LAMP-1 and LAMP-2 are activation markers for normal mast cells (1). They have also been linked to the inflammatory response in that they promote adhesion of human peripheral blood mononuclear cells (PBMC) to vascular endothelium, and therefore possibly the adhesion of PBMC to the site of inflammation (4).

Overview

| Product Name | Anti-LAMP1 Monoclonal Antibody |
|----------------------|--|
| Reactive Species | Hamster, Human, Mouse, Rat |
| Description | Boster Bio Anti-LAMP1 Monoclonal Antibody catalog # M00780-1. Tested in IP, IF, ICC, WB applications. This antibody reacts with Human, Mouse, Rat. |
| Application | IP, IF, ICC, WB |
| Clonality | Monoclonal Ly1C6 |
| Formulation | PBS pH7.4, 50% glycerol, 0.09% sodium azide |
| Storage Instructions | Store at -20°C for one year. Avoid repeated freeze-thaw cycles. |
| Host | Mouse |
| Uniprot ID | P14562 |

Technical Details

| Immunogen | Rat liver lysosomal membrane preparations |
|----------------------------|---|
| Predicted Reactive Species | Canine |
| Cross Reactivity | Detects ~120kDa. |
| Isotype | IgG1 |
| Form | liquid |






888-466-3604 | support@bosterbio.com | www.bosterbio.com

| Concentration | 1 mg/ml |
|---------------------|--|
| Purification | Protein G Purified |
| Suggested Dilutions | Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB (1:1000), ICC/IF (1:1000); optimal dilutions for assays should be determined by the user. |



Anti-LAMP1 Monoclonal Antibody (M00780-1) Images



Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-LAMPCali. San Fran School of Medicine.




Figure 2. Western blot analysis of Lamp1 using anti-Lamp1 antibody (M00780-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Lamp1 antigen affinity purified polyclonal antibody (Catalog # M00780-1) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-Mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # SA1021) with Tanon 5200 system. A specific band was detected for Lamp1.

1 Publications Citing This Product

1. PubMed ID: 33203874, Ma WQ,Sun XJ,Zhu Y,Liu NF. PDK4 promotes vascular calcification by interfering with autophagic activity and metabolic reprogramming. Cell Death Dis.2020 Nov 17;11(11):991.doi:10.1038/s41419-020-03162-w. PMID:33203874;PMCID:PMC7673024.

Visit <u>bosterbio.com/anti-lamp1-antibody-m00780-1-boster.html</u> to see all 1 publications.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-LAMP1 Monoclonal Antibody