

# **Anti-RelB Rabbit Monoclonal Antibody**

Catalog Number: M00836-1

#### **About RELB**

Dynamin-related GTPase required for mitochondrial fusion and regulation of apoptosis. May form a diffusion barrier for proteins stored in mitochondrial cristae. Proteolytic processing in response to intrinsic apoptotic signals may lead to disassembly of OPA1 oligomers and release of the caspase activator cytochrome C (CYCS) into the mitochondrial intermembrane space.

#### Overview

Product Name	Anti-RelB Rabbit Monoclonal Antibody
Reactive Species	Human
Description	Boster Bio Anti-RelB Rabbit Monoclonal Antibody catalog # M00836-1. Tested in WB, IHC, ICC/IF, IP applications. This antibody reacts with Human.
Application	IP, IF, IHC, ICC, WB
Clonality	Monoclonal GCF-18
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q01201

#### **Technical Details**

Immunogen	A synthesized peptide derived from human RelB
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:




# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com


WB 1:5	00-1:1000
IHC 1:5	0-1:200
ICC/IF :	1:50-1:200
IP 1:50	




### Anti-RelB Rabbit Monoclonal Antibody (M00836-1) Images



All lanes use the Antibody at 1:1K dilution for 1 hour at room temperature.



Western blot analysis of RelB expression in Raji cell lysate.



Immunohistochemical analysis of paraffin-embedded human tonsil, using RelB Antibody.

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-RelB Rabbit Monoclonal Antibody