

Anti-Stathmin 1 STMN1 Rabbit Monoclonal Antibody

Catalog Number: M01194

About STMN1

Activated by icilin, eucalyptol, menthol, cold and modulation of intracellular pH. Involved in menthol sensation. Permeable for monovalent cations sodium, potassium, and cesium and divalent cation calcium.

Overview

Product Name	Anti-Stathmin 1 STMN1 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Stathmin 1 STMN1 Rabbit Monoclonal Antibody catalog # M01194. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal DDA-19
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P16949

Technical Details

Immunogen	A synthesized peptide derived from human Stathmin 1
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>WB 1:500-1:2000</p>

	IHC 1:50-1:100 ICC/IF 1:50-1:100 IP 1: 50 FC 1:50
--	---

Anti-Stathmin 1 STMN1 Rabbit Monoclonal Antibody (M01194) Images




Figure 1. Western blot analysis of Stathmin 1 using anti-Stathmin 1 antibody (M00994).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human MDA-MB-453 whole cell lysates,

Lane 2: human SH-SY5Y whole cell lysates,

Lane 3: human Raji whole cell lysates,

Lane 4: rat brain tissue lysates,

Lane 5: rat testis tissue lysates,

Lane 6: rat PC-12 whole cell lysates,

Lane 7: mouse brain tissue lysates,

Lane 8: mouse testis tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5

hour at RT. The membrane was incubated with rabbit anti-

Stathmin 1 antigen affinity purified monoclonal antibody

(Catalog # M00994) at 1:500 overnight at 4°C, then washed

with TBS-0.1%Tween 3 times with 5 minutes each and

probed with a goat anti-rabbit IgG-HRP secondary antibody

at a dilution of 1:500 for 1.5 hour at RT. The signal is

developed using an Enhanced Chemiluminescent detection

(ECL) kit (Catalog # EK1002) with Tanon 5200 system. A

specific band was detected for Stathmin 1 at approximately

17 kDa. The expected band size for Stathmin 1 is at 17 kDa.




Figure 2. IHC analysis of STMN1 using anti-STMN1 antibody (M01194).

STMN1 was detected in a paraffin-embedded section of human colon tissue. Heat mediated antigen retrieval was

performed in EDTA buffer (pH 8.0, epitope retrieval

solution). The tissue section was blocked with 10% goat

serum. The tissue section was then incubated with 1:50

rabbit anti-STMN1 Antibody (M01194) overnight at 4°C.

Peroxidase Conjugated Goat Anti-rabbit IgG was used as

secondary antibody and incubated for 30 minutes at 37°C.

The tissue section was developed using HRP Conjugated

Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with

DAB as the chromogen.




Figure 3. IHC analysis of STMN1 using anti-STMN1 antibody (M01194).

STMN1 was detected in a paraffin-embedded section of human colon tissue. Heat mediated antigen retrieval was

performed in EDTA buffer (pH 8.0, epitope retrieval

solution). The tissue section was blocked with 10% goat

serum. The tissue section was then incubated with 1:50

rabbit anti-STMN1 Antibody (M01194) overnight at 4°C.

Peroxidase Conjugated Goat Anti-rabbit IgG was used as

secondary antibody and incubated for 30 minutes at 37°C.

The tissue section was developed using HRP Conjugated

Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 4. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 5. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of human colon cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 6. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 7. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of human ovarian cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated

Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 8. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of neuroplexus of human colon tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 9. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of neuroplexus of human colon tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 10. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of neuroplexus of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 11. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of neuroplexus of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated

Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 12. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of neuroplexus of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 13. IHC analysis of STMN1 using anti-STMN1 antibody (M01194). STMN1 was detected in a paraffin-embedded section of neuroplexus of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1:50 rabbit anti-STMN1 Antibody (M01194) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Stathmin 1 STMN1 Rabbit Monoclonal Antibody