

# **Anti-CD22 Antibody Picoband™ (monoclonal, 5E7)**

Catalog Number: M01572-2

#### **About CD22**

CD22 is a surface glycoprotein of B lymphocytes that is rapidly phosphorylated on cytoplasmic tyrosines after antigen receptor cross-linking. It is a negative regulator of antigen receptor signaling whose onset of expression at the mature B cell stage may serve to raise the antigen concentration threshold required for B cell triggering. The human CD22 gene is expressed specifically in B lymphocytes and likely has an important function in cell-cell interactions. The B cell coreceptor CD22 plays an important role in regulating signal transduction via the B cell Ag receptor. And CD22 is located within the band region q13.1 of chromosome 19.

#### Overview

Product Name	Anti-CD22 Antibody Picoband™ (monoclonal, 5E7)
Reactive Species	Human
Description	Boster Bio Anti-CD22 Antibody Picoband™ (monoclonal, 5E7) catalog # M01572-2. Tested in Flow Cytometry, WB applications. This antibody reacts with Human.
Application	Flow Cytometry, WB
Clonality	Monoclonal 5E7
Formulation	Each vial contains 4mg Trehalose, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg NaN <sub>3</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Mouse
Uniprot ID	P20273

#### **Technical Details**

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human CD22, different from the related mouse sequence by ten amino acids.
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Mouse IgG (EK1001) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Mouse IgG2b
Form	Lyophilized
Concentration	0
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this



# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide to optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Western blot, 0.1-0.5ug/ml, Human  Flow Cytometry, 1-3ug/1x10 <sup>6</sup> cells, Human	ne
--	----



### Anti-CD22 Antibody Picoband™ (monoclonal, 5E7) (M01572-2) Images




Figure 1. Western blot analysis of CD22 using anti ZO-1 antibody (M01572-2).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: human Raji tissue lysates,

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with mouse anti-CD22 antigen affinity purified polyclonal antibody (Catalog # M01572-2) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-mouse IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1001) with Tanon 5200 system. A specific band was detected for CD22 at approximately 140KD. The expected band size for CD22is at 95KD.




Figure 2. Flow Cytometry analysis of PBMC cells using anti-CD22 antibody (M01572-2).

Overlay histogram showing PBMC cells stained with M01572-2 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with mouse anti-CD22 Antibody (M01572-2,  $1ug/1x10^6$  cells) for 30 min at  $20^{\circ}$ C. DyLight® 488 conjugated goat anti-mouse IgG (BA1126, 5- $10ug/1x10^6$  cells) was used as secondary antibody for 30 minutes at  $20^{\circ}$ C. Isotype control antibody (Green line) was mouse IgG ( $1ug/1x10^6$ ) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-CD22 Antibody (monoclonal, 5E7)