

Anti-67kDa Laminin Receptor RPSA Rabbit Monoclonal Antibody

Catalog Number: M01691

About RPSA

Furin is likely to represent the ubiquitous endoprotease activity within constitutive secretory pathways and capable of cleavage at the RX (K/R) R consensus motif.

Overview

Product Name	Anti-67kDa Laminin Receptor RPSA Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-67kDa Laminin Receptor RPSA Rabbit Monoclonal Antibody catalog # M01691. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal AOA0-18
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P08865

Technical Details

Immunogen	A synthesized peptide derived from human 67kDa Laminin Receptor
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used:

	WB 1:500-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200 IP 1:50 FC 1:50
--	--

Anti-67kDa Laminin Receptor RPSA Rabbit Monoclonal Antibody (M01691) Images






Figure 1. Western blot analysis of 67kDa Laminin Receptor using anti-67kDa Laminin Receptor antibody (M01691). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HeLa whole cell lysates,
Lane 2: human K562 whole cell lysates,
Lane 3: human HepG2 whole cell lysates,
Lane 4: human COLO320 whole cell lysates,
Lane 5: mouse liver tissue lysates,
Lane 6: mouse NIH/3T3 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-67kDa Laminin Receptor antigen affinity purified monoclonal antibody (Catalog # M01691) at 1:500 overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:500 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for 67kDa Laminin Receptor at approximately 40 kDa. The expected band size for 67kDa Laminin Receptor is at 33 kDa.



All lanes use the Antibody at 1:6K dilution for 1 hour at room temperature.



Western blot analysis of 67kDa Laminin Receptor expression in (1) HeLa cell lysate; (2) NIH/3T3 cell lysate; (3) PC-12 cell lysate.

1 Publications Citing This Product

neural stem cells by sustained release of Shh from TG2 gene-modified EMSC co-culture in vitro. Amino Acids. 2020 Nov 27. doi: 10.1007/

Visit bosterbio.com/anti-67kda-laminin-receptor-rabbit-monoclonal-antibody-m01691-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-67kDa Laminin Receptor RPSA Rabbit Monoclonal Antibody