

Anti-UCHL3 Rabbit Monoclonal Antibody

Catalog Number: M05004-1

About UCHL3

Putative transcription factor involved in pancreas development and function.

Overview

Product Name	Anti-UCHL3 Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-UCHL3 Rabbit Monoclonal Antibody catalog # M05004-1. Tested in WB, IHC applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Monoclonal 26U65
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P15374

Technical Details

Immunogen	A synthesized peptide derived from UCHL3
Isotype	IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB 1:500-1:2000 IHC 1:50-1:200

Anti-UCHL3 Rabbit Monoclonal Antibody (M05004-1) Images




Figure 1. Western blot analysis of UCHL3 using anti-UCHL3 antibody (M05004-1).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human K562 whole cell lysates,

Lane 2: human 293T whole cell lysates,

Lane 3: human A431 whole cell lysates,

Lane 4: human CACO-2 whole cell lysates,

Lane 5: rat heart tissue lysates,

Lane 6: mouse heart tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5

hour at RT. The membrane was incubated with rabbit anti-UCHL3 antigen affinity purified monoclonal antibody (Catalog # M05004-1) at 1:500 overnight at 4°C, then

washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:1000 for 1.5 hour at RT. The

signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200

system. A specific band was detected for UCHL3 at approximately 26 kDa. The expected band size for UCHL3 is at 26 kDa.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-UCHL3 Rabbit Monoclonal Antibody