

Anti-beta II Tubulin Rabbit Monoclonal Antibody

Catalog Number: M09008

About TUBB4B

Putative transcription factor involved in pancreas development and function.

Overview


Product Name	Anti-beta II Tubulin Rabbit Monoclonal Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-beta II Tubulin Rabbit Monoclonal Antibody catalog # M09008. Tested in WB, IHC, ICC/IF, IP, Flow Cytometry applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IP, IF, IHC, ICC, WB
Clonality	Monoclonal DOF-20
Formulation	Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol, 0.4-0.5mg/ml BSA.
Storage Instructions	Store at -20°C for one year. For short term storage and frequent use, store at 4°C for up to one month. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P68371/Q13885/Q9BVA1

Technical Details


Immunogen	A synthesized peptide derived from human beta II Tubulin
Isotype	Rabbit IgG
Form	Liquid
Concentration	Actual concentration vary by lot. Use suggested dilution ratio to decide dilution procedure.
Purification	Affinity-chromatography
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: WB 1:1000-1:2000 IHC 1:50-1:200 ICC/IF 1:50-1:200

	IP 1:50 FC 1:50
--	--------------------


Anti-beta II Tubulin Rabbit Monoclonal Antibody (M09008) Images




All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.





All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Western blot analysis of beta II Tubulin expression in C6 cell lysate.

All lanes use the Antibody at 1:1W dilution for 1 hour at room temperature.



Immunohistochemical analysis of paraffin-embedded mouse brain, using beta II Tubulin Antibody.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-beta II Tubulin Rabbit Monoclonal Antibody