

# Anti-Acetyl-Histone H4 (Lys8) HIST1H4A Rabbit Monoclonal Antibody, Clone#RM201

Catalog Number: M14495-7

### Overview

Product Name	Anti-Acetyl-Histone H4 (Lys8) HIST1H4A Rabbit Monoclonal Antibody, Clone#RM201
Reactive Species	Human, Vertebrates
Description	Boster Bio Anti-Acetyl-Histone H4 (Lys8) HIST1H4A Rabbit Monoclonal Antibody, Clone#RM201 (Catalog # M14495-7). Tested in ChIP, WB, ELISA, Luminex, ICC applications. This antibody reacts with Human, Vertebrates.
Application	ChIP, ELISA, ICC, WB, Luminex
Clonality	Monoclonal RM201
Formulation	50% Glycerol/PBS with 1% BSA and 0.09% sodium azide
Storage Instructions	Store at -20°C for one year. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P62805

#### **Technical Details**

Immunogen	An acetyl-peptide corresponding to the Acetyl-Histone H4 (Lys8).
Cross Reactivity	This antibody reacts to Histone H4 acetylated at Lysine 8 (K8ac). No cross-reactivity with other acetylated Lysines in Histone H4.
Isotype	Rabbit IgG
Form	Liquid
Concentration	1 mg/mL
Purification	Protein A affinity purified from an animal origin-free culture supernatant
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  ChIP: 1µg-5 µg  WB: 0.5 ug/mL - 2 ug/mL  ELISA: 0.2 ug/mL - 1 ug/mL  Luminex: 0.05 ug/mL - 1 ug/mL



### BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

ICC: 0.5 ug/mL- 2 ug/mL.



## Anti-Acetyl-Histone H4 (Lys8) HIST1H4A Rabbit Monoclonal Antibody, Clone#RM201 (M14495-7) Images




Figure 1. Western Blotting result Western Blot of (1) acid extracts of HeLa cells treated with sodium butyrate, (2) acid extracts of HeLa cells untreated, and (3) recombinant Histone H4. Using RM201 at 0.5 ug/mL, showed a band of Histone H4 acetylated at Lysine 8 in HeLa cells




Figure 2. Specificity Test result RM201 specifically reacts to Histone H4 acetylated at Lysine 8 (K8ac). No cross reactivity with unmodified Lysine 8 (K8 ctrl), acetylated Lysine 5 (K5ac), Lysine 12 (K12ac), Lysine 16 (K16ac), Lysine 20 (K20ac), Lysine 31 (K31ac), or Lysine 91 (K91) in Histone H4.




Figure 3. ICC result Immunocytochemistry of HeLa cells treated with sodium butyrate, using Acetyl-Histone H4 (Lys8) Rabbit Monoclonal Antibody (Clone RM201) (red). Actin filaments have been labeled with fluorescein phalloidin (green).

### Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.