

Anti-Fibroblast growth factor 4 FGF4 Antibody

Catalog Number: PA1033

About FGF4

Fibroblast growth factor 4 (FGF4), also known as Heparin Secretary Transforming (HSTF1). HST1, for which the designation HSTF1 was proposed for human gene nomenclature, is a heparin-binding growth factor with significant homology to human fibroblast growth factors and the mouse Int-2 protein. By in situ hybridization, Adelaide et al. (1988) mapped the HST gene to chromosome 11q13. The HST1 protein is a heparin-binding growth factor with significant homology with human fibroblast growth factors and the mouse Int-2 protein.

Overview

Product Name	Anti-Fibroblast growth factor 4 FGF4 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Fibroblast growth factor 4 FGF4 Antibody catalog # PA1033. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P08620

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human FGF4, identical to the related rat and mouse sequences.
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.




BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, Rat, Mouse



Anti-Fibroblast growth factor 4 FGF4 Antibody (PA1033) Images



Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Fibroblast growth factor 4 FGF4 Antibody