

Anti-Lamin B1/LMNB1 Antibody

Catalog Number: PA1048

About LMNB1

Lamin-B1 is a protein that in humans is encoded by the LMNB1 gene. The nuclear lamina consists of a two-dimensional matrix of proteins located next to the inner nuclear membrane. The lamin family of proteins make up the matrix and are highly conserved in evolution. During mitosis, the lamina matrix is reversibly disassembled as the lamin proteins are phosphorylated. Lamin proteins are thought to be involved in nuclear stability, chromatin structure and gene expression. Vertebrate lamins consist of two types, A and B. This gene encodes one of the two B type proteins, B1.

Overview

Product Name	Anti-Lamin B1/LMNB1 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Lamin B1/LMNB1 Antibody catalog # PA1048. Tested in IF, IHC, IHC-F, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IF, IHC, IHC-F, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P20700

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Lamin B1, different from the related rat sequence by one amino acid, and from the related mouse sequence by three amino acids.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P), IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Rat, By Heat Immunohistochemistry (Frozen Section), 0.5-1ug/ml, Rat Immunocytochemistry , 0.5- 1ug/ml, Human Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human

Anti-Lamin B1/LMNB1 Antibody (PA1048) Images




Figure 1. Western blot analysis of Lamin B1 using anti-Lamin B1 antibody (PA1048).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Daudi whole cell lysates,

Lane 2: human THP-1 whole cell lysates,

Lane 3: human HL-60 whole cell lysates,

Lane 4: rat lung tissue lysates,

Lane 5: mouse lung tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Lamin B1 antigen affinity purified polyclonal antibody (Catalog # PA1048) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Lamin B1 at approximately 70 kDa. The expected band size for Lamin B1 is at 70 kDa.




Figure 2. IHC analysis of Lamin B1 using anti-Lamin B1 antibody (PA1048).

Lamin B1 was detected in a paraffin-embedded section of Human Mammary Cancer tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-Lamin B1 Antibody (PA1048) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.





Figure 3. IHC analysis of Lamin B1 using anti-Lamin B1 antibody (PA1048).

Lamin B1 was detected in a frozen section of Rat Intestine tissue. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 ug/ml rabbit anti-Lamin B1 Antibody (PA1048) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC) (Catalog # SA1022) with DAB as the chromogen.

Figure 4. ICC analysis of Lamin B1 using anti-Lamin B1



antibody (PA1048). Lamin B1 was detected in an immunocytochemical section of A549 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 1 ug/ml rabbit anti-Lamin B1 Antibody (PA1048) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Figure 5. IF analysis of Lamin B1 and Tubulin beta using anti-Lamin B1 antibody (PA1048) and anti-Tubulin beta antibody (M01857-3). Lamin B1 and Tubulin beta were detected in immunocytochemical section of SiHa cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5ug/mL rabbit anti-Lamin B1 antibody (PA1048) and mouse anti-Tubulin beta Antibody (M01857-3) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) and DyLight®488 Conjugated Goat Anti-Mouse IgG (BA1126) were used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. IF analysis of Lamin B1 using anti-Lamin B1 antibody (PA1048). Lamin B1 was detected in an immunocytochemical section of SiHa cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins. The cells were blocked with 10% goat serum. And then incubated with 5 ug/mL rabbit anti-Lamin B1 Antibody (PA1048) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:100 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.

24 Publications Citing This Product

1. PubMed ID: 33125154, Huang T,Luo X,Wu B,Peng P,Dai Y,Hu G,Qiu H,Yuan X.Pyrotinib enhances the radiosensitivity of HER2⁺overexpressing gastric and breast cancer cells.Oncol Rep.2020 Dec;44(6):2634-2644.doi:10.3892/or.2020.7820.Epub 2020 Oct 21.PMID:33125154;PMCID:PMC7640366.
2. PubMed ID: -, Na,L.,Q.B.,Xiumei,Z.et al.Research into the intervention effect of folic acid on arsenic-induced heart abnormalities in fetal rats during the periconception period.BMC Cardiovasc Disord 20, 139 (2020).https://doi.org/10.1186/s12872-020-01418-z
3. PubMed ID: 32848589, Zhang B,Zhong Q,Chen X,Wu X,Sha R,Song G,Zhang C,Chen X.Neuroprotective Effects of Celastrol on Transient Global Cerebral Ischemia Rats via Regulating HMGB1/NF-kappaB Signaling Pathway.Front Neurosci.2020 Aug 11;14:847.doi:10.3389/fnins.2020.00847.PMID:328485

Visit bosterbio.com/anti-lamin-b1-antibody-pa1048-boster.html to see all 24 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Lamin B1/LMNB1 Antibody