

## Anti-Myeloperoxidase/MPO Antibody

Catalog Number: PA1054

### About MPO

Myeloperoxidase (MPO) is a mammalian phagocyte hemoprotein though to primarily mediate host defense reactions. It is abundantly expressed in neutrophils and secreted during their activation. Myeloperoxidase is part of the host defense system of human polymorphonuclear leukocytes, responsible for microbicidal activity against a wide range of organisms. It is located in the nucleus as well as in the cytoplasm. Intranuclear MPO may help to protect DNA against damage resulting from oxygen radicals produced during myeloid cell maturation and function.

### Overview

|                      |                                                                                                                                                                                                       |
|----------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Product Name         | Anti-Myeloperoxidase/MPO Antibody                                                                                                                                                                     |
| Reactive Species     | Human, Mouse, Rat                                                                                                                                                                                     |
| Description          | Boster Bio Anti-Myeloperoxidase/MPO Antibody catalog # PA1054. Tested in Flow Cytometry, IF, IHC, WB applications. This antibody reacts with Human, Mouse, Rat.                                       |
| Application          | Flow Cytometry, IF, IHC, WB                                                                                                                                                                           |
| Clonality            | Polyclonal                                                                                                                                                                                            |
| Formulation          | Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na <sub>2</sub> HPO <sub>4</sub> , 0.05mg Thimerosal, 0.05mg NaN <sub>3</sub> .                                                                         |
| Storage Instructions | Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles. |
| Host                 | Rabbit                                                                                                                                                                                                |
| Uniprot ID           | P05164                                                                                                                                                                                                |

### Technical Details

|                               |                                                                                                                                                                                  |
|-------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Immunogen                     | A synthetic peptide corresponding to a sequence at the C-terminus of human MPO, different from the related mouse and rat sequences by one amino acid.                            |
| Predicted Reactive Species    | Chicken                                                                                                                                                                          |
| Recommended Detection Systems | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P). |
| Cross Reactivity              | No cross-reactivity with other proteins                                                                                                                                          |
| Isotype                       | Rabbit IgG                                                                                                                                                                       |
| Form                          | Lyophilized                                                                                                                                                                      |
| Concentration                 | Adding 0.2 ml of distilled water will yield a concentration of 500 µg/ml.                                                                                                        |

|                     |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|---------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purification        | Immunogen affinity purified.                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                         |
| Suggested Dilutions | <p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5µg/ml, Human, Mouse, Rat</p> <p>Immunohistochemistry (Paraffin-embedded Section), 2-5µg/ml, Human, Mouse, Rat, By Heat</p> <p>Immunofluorescence, 5 ug/ml, Human</p> <p>Flow Cytometry, 1-3 ug/1x10<sup>6</sup> cells, Human</p> |

## Anti-Myeloperoxidase/MPO Antibody (PA1054) Images




Figure 1. Western blot analysis of MPO using anti-MPO antibody (PA1054).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HL-60 whole cell lysates,

Lane 2: rat thymus tissue lysates,

Lane 3: mouse spleen tissue lysates,

Lane 4: mouse thymus tissue lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-MPO antigen affinity purified polyclonal antibody (Catalog # PA1054) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for MPO at approximately 60 kDa. The expected band size for MPO is at 84 kDa.




Figure 2. IHC analysis of MPO using anti-MPO antibody (PA1054).

MPO was detected in a paraffin-embedded section of human tonsil tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MPO Antibody (PA1054) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.





Figure 3. IHC analysis of MPO using anti-MPO antibody (PA1054).

MPO was detected in a paraffin-embedded section of mouse spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MPO Antibody (PA1054) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Figure 4. IHC analysis of MPO using anti-MPO antibody



(PA1054).  
MPO was detected in a paraffin-embedded section of rat lymphaden tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-MPO Antibody (PA1054) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.




Figure 5. IF analysis of MPO using anti-MPO antibody (PA1054).  
MPO was detected in a paraffin-embedded section of human spleen tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-MPO Antibody (PA1054) overnight at 4°C. DyLight488 Conjugated Goat Anti-Rabbit IgG (BA1127) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 6. IF analysis of MPO using anti-MPO antibody (PA1054).  
MPO was detected in a paraffin-embedded section of human appendicitis tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 5 ug/ml rabbit anti-MPO Antibody (PA1054) overnight at 4°C. Cy3 Conjugated Goat Anti-Rabbit IgG (BA1032) was used as secondary antibody at 1:500 dilution and incubated for 30 minutes at 37°C. The section was counterstained with DAPI. Visualize using a fluorescence microscope and filter sets appropriate for the label used.




Figure 7. Flow Cytometry analysis of HL-60 cells using anti-MPO antibody (PA1054).  
Overlay histogram showing HL-60 cells stained with PA1054 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-MPO Antibody (PA1054, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

## 6 Publications Citing This Product

1. PubMed ID: -, Wei Gao,Qiushi Huang,Ting Qin et al.HSP60 regulates monosodium urate crystal-induced inflammation by activating the TLR4-NF-

kappa B-MyD88 signaling pathway and disrupting mitochondrial function, 30 April 2020, PREPRINT (Version 1) available at Research Square

2. PubMed ID: 27557503, Protective effects of Huangqin Decoction against ulcerative colitis and associated cancer in mice

3. PubMed ID: 28765694, Lactobacillus acidophilus alleviates pouchitis after ileal pouch-anal anastomosis in rats

Visit [bosterbio.com/anti-mpo-antibody-pa1054-boster.html](https://bosterbio.com/anti-mpo-antibody-pa1054-boster.html) to see all 6 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Myeloperoxidase/MPO Antibody