

Anti-NF-kB p65/RELA Antibody

Catalog Number: PA1125

About RELA

The p65 (RELA) heterodimer is the most abundant form of NFkB. This gene is located on 11q13, which consists of 10 exons and spans about 8.1 kb of DNA. In rat sciatic nerves, the expression of the activated p65 subunit of NFkB was high in the nuclei of premyelinating Schwann cells and then progressively declined until it was nearly absent in adults. The transcriptional activity of NF-kappa-B is stimulated upon phosphorylation of its p65 subunit on serine-276 by protein kinase A (PKA). The transcriptional coactivator CBP/p300 associates with NF-kappa-B p65 through 2 sites, an N-terminal domain that interacts with the C-terminal region of unphosphorylated p65, and a second domain that only interacts with p65 phosphorylated on serine-276.

Overview


Product Name	Anti-NF-kB p65/RELA Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-NF-kB p65/RELA Antibody catalog # PA1125. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q04206

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human NF-kB p65, identical to the related rat sequence, and different from the related mouse sequence by one amino acid.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p>

Anti-NF-kB p65/RELA Antibody (PA1125) Images



Anti-NF-kB p65 antibody, PA1125, Western blotting
All lanes: Anti NF-kB p65 (PA1125) at 0.5ug/ml
Lane 1: Human Colon Cancer Tissue Lysate at 50ug
Lane 2: HELA Whole Cell Lysate at 40ug
Predicted bind size: 65KD
Observed bind size: 65KD

49 Publications Citing This Product

1. PubMed ID: 33607154, Liao L,Huang L,Wei X,Yin L,Wei X,Li T.Bioinformatic and biochemical studies of formononetin against liver injure.Life Sci.2021 Feb 16:119229.doi:10.1016/j.lfs.2021.119229.Epub ahead of print.PMID:33607154.
2. PubMed ID: 33607154, Liao L,Huang L,Wei X,Yin L,Wei X,Li T.Bioinformatic and biochemical studies of formononetin against liver injure.Life Sci.2021 Feb 16:119229.doi:10.1016/j.lfs.2021.119229.Epub ahead of print.PMID:33607154.
3. PubMed ID: 32855700, Liu L,Liu F,Sun Z,Peng Z,You T,Yu Z. LncRNA NEAT1 promotes apoptosis and inflammation in LPS-induced sepsis models by targeting miR-590-3p. Exp Ther Med.2020 Oct; 20(4):3290-3300. doi:10.3892/etm.2020.9079. Epub 2020 Jul 29.PMID:32855700;PMCID:PMC7444425.

Visit bosterbio.com/anti-nf-kb-p65-antibody-pa1125-boster.html to see all 49 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-NF-kB p65/RELA Antibody