

Anti-ERK1/MAPK3 Antibody

Catalog Number: PA1343

About MAPK3

Mitogen-activated protein kinase 3 is an enzyme that in humans is encoded by the MAPK3 gene. The protein encoded by this gene is a member of the MAP kinase family. MAP kinases, also known as extracellular signal-regulated kinases (ERKs), act in a signaling cascade that regulates various cellular processes such as proliferation, differentiation, and cell cycle progression in response to a variety of extracellular signals. This kinase is activated by upstream kinases, resulting in its translocation to the nucleus where it phosphorylates nuclear targets. Alternatively spliced transcript variants encoding different protein isoforms have been described. MAPK3 gene is mapped to human chromosome 16 by hybrid cell panel analysis.

Overview


Product Name	Anti-ERK1/MAPK3 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-ERK1/MAPK3 Antibody catalog # PA1343. Tested in IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P27361

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human ERK1, identical to the related mouse and rat sequences.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.


Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Rat, Mouse, By Heat</p> <p>Immunocytochemistry , 0.5-1ug/ml, Human, Mouse, Rat</p> <p>Western blot, 0.1-0.5ug/ml, Human, Rat, Mouse</p>

Anti-ERK1/MAPK3 Antibody (PA1343) Images




Anti-ERK1 antibody, PA1343, Western blotting

Lane 1: Rat Spleen Tissue Lysate
Lane 2: Rat Thymus Tissue Lysate
Lane 3: Rat Skeletal Muscle Tissue Lysate
Lane 4: Rat Kidney Tissue Lysate
Lane 5: HELA Cell Lysate
Lane 6: JURKAT Cell Lysate
Lane 7: RAJI Cell Lysate
Lane 8: COLO320 Cell Lysate



Anti-ERK1 antibody, PA1343, IHC(P)

IHC(P): Human Mammary Cancer Tissue



Anti-ERK1 antibody, PA1343, IHC(P)

ICC: HELA Cell




Figure 4. IHC analysis of ERK1 using anti-ERK1 antibody (PA1343).

ERK1 was detected in paraffin-embedded section of rat intestine tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-ERK1 Antibody (PA1343) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

10 Publications Citing This Product

1. PubMed ID: 17392578, Hu Cp, Feng Jt, Tang Yl, Zhu Jq, Lin Mj, Yu Me. Mediators Inflamm. 2006;2006(5):84829. Lif Upregulates Expression Of Nk-1R In Nhbe Cells.

2. PubMed ID: 24809625, Yang Sl, Han R, Liu Y, Hu Ly, Li Xl, Zhu Ly. Wound Repair Regen. 2014 Jul-Aug;22(4):548-54. Doi: 10.1111/Wrr.12195.

Negative Pressure Wound Therapy Is Associated With Up-Regulation Of Bfgf And Erk1/2 In Human Diabetic Foot Wounds.

3. PubMed ID: 25337216, Liu Y, Zhu Z, Xiong Z, Zheng J, Hu Z, Qiu J. Int J Clin Exp Pathol. 2014 Aug 15;7(9):5750-61. Ecollection 2014. Knockdown Of Protein Tyrosine Phosphatase Receptor U Inhibits Growth And Motility Of Gastric Cancer Cells.

Visit bosterbio.com/anti-erk1-antibody-pa1343-boster.html to see all 10 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-ERK1/MAPK3 Antibody