

Anti-HSD17B6 Antibody

Catalog Number: PA1386

About HSD17B6

Hydroxysteroid 17-beta dehydrogenase 6 is an enzyme that in humans is encoded by the HSD17B6 gene. The protein encoded by this gene has both oxidoreductase and epimerase activities and is involved in androgen catabolism. Baker ME et al point out expression of 17beta-HSDs had an important role in the early evolution of the physiological response to androgens and estrogens. Biswas and Russell concluded that 17beta-HSD6 and RoDH play opposing roles in androgen action; 17beta-HSD6 inactivates 3alpha-adial by conversion to androsterone and RoDH activates 3alpha-adial by conversion to dihydrotestosterone. The synthesis of an active steroid hormone by back conversion of an inactive metabolite represents a potentially important mechanism by which the steady state level of a transcriptional effector can be regulated.

Overview

Product Name	Anti-HSD17B6 Antibody
Reactive Species	Human
Description	Boster Bio Anti-HSD17B6 Antibody catalog # PA1386. Tested in IHC, WB applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O14756

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human HSD17B6.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat</p> <p>Western blot, 0.1-0.5ug/ml, Human</p>

Anti-HSD17B6 Antibody (PA1386) Images



Anti-HSD17B6 antibody, PA1386, Western blotting
All lanes: Anti HSD17B6 (PA1386) at 0.5ug/ml
Lane 1: Human Placenta Tissue Lysate at 50ug
Lane 2: MCF-7 Whole Cell Lysate at 40ug
Predicted bind size: 35KD
Observed bind size: 35KD




Figure 2. IHC analysis of HSD17B6 using anti-HSD17B6 antibody (PA1386).

HSD17B6 was detected in paraffin-embedded section of human liver cancer tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-HSD17B6 Antibody (PA1386) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HSD17B6 Antibody