

Anti-Fractalkine CX3CL1 Antibody

Catalog Number: PA1400

About CX3CL1

Chemokine (C-X3-C motif) ligand 1, also known as CX3CL1, is a large cytokine protein. It is said to be the only known member of the CX3C chemokine family. It is also commonly known under the names fractalkine (in humans) and neurotactin (in mice). Its gene is mapped to the long arm of chromosome 8. Being the large protein, it has got 373 amino acids. Imai et al. (1997) identified a 7-transmembrane high-affinity receptor (CX3CR1; 601470) for fractalkine and showed that it mediates both the adhesive and migratory functions of fractalkine.

Overview

Product Name	Anti-Fractalkine CX3CL1 Antibody
Reactive Species	Human
Description	Boster Bio Anti-Fractalkine CX3CL1 Antibody catalog # PA1400. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P78423

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human CX3CL1.
Predicted Reactive Species	Monkey
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.

Suggested Dilutions

Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.


If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.

Some PubMed article(s) citing the expression level of this target are as follows:

Boster Bio's internal QC testing used:

Western blot, 0.1-0.5ug/ml, Human

Anti-Fractalkine CX3CL1 Antibody (PA1400) Images



Anti-CX3CL1 antibody, PA1400, Western blotting
Lane 1: Recombinant Human Fractalkine Protein 10ng
Lane 2: Recombinant Human Fractalkine Protein 5ng
Lane 3: Recombinant Human Fractalkine Protein 2.5ng

3 Publications Citing This Product

1. PubMed ID: 10.1002/ar.22430, Distribution Change of Mast Cells in Human Nasal Polyps

2. PubMed ID: 10.1002/ar.22430, Distribution Change of Mast Cells in Human Nasal Polyps

3. PubMed ID: 10.1002/ar.22430, Distribution Change of Mast Cells in Human Nasal Polyps

Visit bosterbio.com/anti-fractalkine-cx3cl1-antibody-pa1400-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Fractalkine CX3CL1 Antibody