

Anti-Thrombospondin 2/THBS2 Antibody

Catalog Number: PA1417

About THBS2

Thrombospondin-2 (THBS2) is a protein that in humans is encoded by the THBS2 gene. The protein encoded by this gene belongs to the thrombospondin family. The THBS2 is mapped to 6q27 and it is located on chromosome 17. The gene was transcribed in fibroblasts, smooth muscle cells, and an osteosarcoma cell line. It functions as a protein inhibitor of tumor growth and angiogenesis and modulates the cell surface properties of mesenchymal cells and be involved in cell adhesion and migration.

Overview

Product Name	Anti-Thrombospondin 2/THBS2 Antibody
Reactive Species	Human
Description	Boster Bio Anti-Thrombospondin 2/THBS2 Antibody catalog # PA1417. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.01mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P35442

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human Thrombospondin 2, different from the mouse sequence by two amino acids.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human



Anti-Thrombospondin 2/THBS2 Antibody (PA1417) Images




Figure 1. Western blot analysis of THBS2 using anti-THBS2 antibody (PA1417).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human MCF-7 whole cell lysates,

Lane 3: human U87 whole cell lysates,

Lane 4: human U251 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-THBS2 antigen affinity purified polyclonal antibody (Catalog # PA1417) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for THBS2 at approximately 160 kDa. The expected band size for THBS2 is at 130 kDa.

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Thrombospondin 2/THBS2 Antibody