

# **Anti-CAMKK2 Antibody**

Catalog Number: PA1497

#### **About CAMKK2**

CAMKK2, Calcium/calmodulin-dependent protein kinase kinase 2, is an enzyme that in humans is encoded by the CAMKK2 gene. The product of this gene belongs to the serine/threonine-specific protein kinase family, and to the Ca++/calmodulin-dependent protein kinase subfamily. This protein plays a role in the calcium/calmodulin-dependent (CaM) kinase cascade by phosphorylating the downstream kinases CaMK1 and CaMK4. CaMKK2 regulates production of the appetite stimulating hormone neuropeptide Y and functions as an AMPK kinase in the hypothalamus. It also has an important role in the development of hyperalgesia and tolerance to opioid analgesic drugs, through reduction in downstream signalling pathways and mu opioid receptor downregulation. Inhibition of CaMKK2 in mice reduces appetite and promotes weight loss.

#### Overview

Product Name	Anti-CAMKK2 Antibody
Reactive Species	Mouse, Rat
Description	Boster Bio Anti-CAMKK2 Antibody catalog # PA1497. Tested in IHC, WB applications. This antibody reacts with Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q96RR4

#### **Technical Details**

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human CAMKK2, identical to the related rat and mouse sequences.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized





<b>BOSTER</b>
antibody and ELISA experts

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Rat, By Heat Western blot, 0.1-0.5ug/ml, Mouse, Rat




## Anti-CAMKK2 Antibody (PA1497) Images

130KD - 1 2 3 4 100KD -70KD -55KD -35KD -25KD -

Anti-CAMKK2 antibody, PA1497, Western blotting

Lane 1: Rat Brain Tissue Lysate Lane 2: Rat Brain Tissue Lysate Lane 3: Mouse Brain Tissue Lysate Lane 4: Mouse Brain Tissue Lysate



Anti-CAMKK2 antibody, PA1497, IHC(P) IHC(P): Rat Brain Tissue

### Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.





Anti-CAMKK2 Antibody