

Anti-HSD17B2 Antibody

Catalog Number: PA1604

About HSD17B2

HSD17B2 (17-BETA-HYDROXYSTEROID DEHYDROGENASE II), also called 17-BETA-HSD II, is an enzyme which has 387 amino acids with a predicted molecular weight of 42,782 and associates with the membranes of the endoplasmic reticulum. Its cytogenetic location is 16q23.3. The type 2 enzyme was capable of catalyzing the interconversion of testosterone and androstenedione, as well as estradiol and estrone. HSD17B2 mRNA was detected in 18 of 42 (43%) adenomas but not in prolactinomas. In the human endometrium, inactivation of 17-beta-estradiol to estrone is catalyzed by HSD17B2. And HSD17B2 activity distinguishes between disease-free and diseased endometria. HSD17B2 efficiently catalyzes the oxidative metabolism of androgens and estrogens, and it is expressed in a large series of human peripheral tissues. The previous paradigm that HSD17B2 activity in the endometrium is elevated during the secretory phase is confined to diseased endometrium but not to disease-free endometrium.

Overview


Product Name	Anti-HSD17B2 Antibody
Reactive Species	Human
Description	Boster Bio Anti-HSD17B2 Antibody catalog # PA1604. Tested in IHC, WB applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P37059

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human HSD17B2.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and IHC(F).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Immunohistochemistry (Frozen Section), 0.5-1ug/ml, Human, -</p> <p>Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat</p> <p>Western blot, 0.1-0.5ug/ml, Human</p>


Anti-HSD17B2 Antibody (PA1604) Images



Anti-HSD17B2 antibody, PA1604, Western blotting
All lanes: Anti HSD17B2 (PA1604) at 0.5ug/ml
WB: Human Placenta Tissue Lysate at 50ug
Predicted bind size: 43KD
Observed bind size: 43KD



Anti-HSD17B2 antibody, PA1604, IHC(P)
IHC(P): Human Placenta Tissue



Anti-HSD17B2 antibody, PA1604, IHC(F)
IHC(F): Human Placenta Tissue

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-HSD17B2 Antibody