

# **Anti-MyD88 Antibody**

Catalog Number: PA1660

#### **About MYD88**

MYD88 (MYELOID DIFFERENTIATION PRIMARY RESPONSE GENE 88), is a protein that, in humans, is encoded by the MYD88 gene. MyD88 is a key downstream adapter for most Toll-like receptors (TLRs) and interleukin-1 receptors (IL1Rs). And it is mapped on 3p22.2. MYD88 encodes a cytosolic adapter protein that plays a central role in the innate and adaptive immune response. This protein functions as an essential signal transducer in the interleukin-1 and Toll-like receptor signaling pathways. Qverexpression of MYD88 caused an increase in the level of transcription from the interleukin-8 promoter. The C-terminal domain of MYD88 has significant sequence similarity to the cytoplasmic domain of IL1RAP. Inhibiting the IL1R-MYD88 pathway in vivo could block the damage from acute inflammation that occurs in response to sterile cell death, and do so in a way that might not compromise tissue repair or host defense against pathogens.

#### Overview

Product Name	Anti-MyD88 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-MyD88 Antibody catalog # PA1660. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.01 mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q99836

#### **Technical Details**

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human MyD88, different from the related rat and mouse sequences by one amino acid.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG






888-466-3604 | support@bosterbio.com | www.bosterbio.com

Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, Rat, Mouse, By Heat Western blot, 0.1-0.5ug/ml, Human, Rat, Mouse




### Anti-MyD88 Antibody (PA1660) Images



Anti-MyD88 antibody, PA1660, Western blotting

Lane 1: Rat Spleen Tissue Lysate Lane 2: Rat Thymus Tissue Lysate

Lane 3: JURKAT Cell Lysate Lane 4: RAJI Cell Lysate



Anti-MyD88 antibody, PA1660, IHC(P) IHC(P): Human Tonsil Tissue

### **5 Publications Citing This Product**

1. PubMed ID: 32593156, Li X,Shi MQ,Chen C,Du JR.Phthalide derivative CD21 ameliorates ischemic brain injury in a mouse model of global cerebral ischemia: involvement of inhibition of NLRP3.Int Immunopharmacol.2020 Sep;86:106714.doi: 10.1016/j.intimp.2020.106714.Epub 2020 Jun 24

- 2. PubMed ID: 26468333, High glucose induces and activates Toll-like receptor 4 in endothelial cells of diabetic retinopathy
- 3. PubMed ID: 25299052, Toll-Like Receptor 4 Prompts Human Breast Cancer Cells Invasiveness via Lipopolysaccharide Stimulation and Is Overexpressed in Patients with Lymph Node Metastasis

Visit <u>bosterbio.com/anti-myd88-antibody-pa1660-boster.html</u> to see all 5 publications.

## Submit a product review to Biocompare.com







Anti-MyD88 Antibody