

# **Anti-Interleukin-3 IL3 Antibody**

Catalog Number: PA1748

### **About IL3**

The gene IL-3 encodes interleukin 3, a hematopoietic colony-stimulating factor (CSF) that is capable of supporting the proliferation of a broad range of hematopoietic cell types. Interleukin-3 (IL-3), a protein of 140 amino acids, is chemically synthesized by means of an automated peptide synthesizer and is shown to have the biological activities attributed to native IL-3. The cDNA sequence for murine interleukin-3, one of the colony stimulating factors that regulate haematopoiesis, codes for a polypeptide of 166 amino acids including a putative signal peptide. The mouse IL 3 gene is located on chromosome 11. The human gene encoding IL 3 is tandemly arrayed on the long arm of chromosome 5.5 The standard product used in this kit is recombinant human IL-3, consisting of 133 amino acids with the molecular mass of 15KDa.

### Overview

Product Name	Anti-Interleukin-3 IL3 Antibody
Reactive Species	Human
Description	Boster Bio Anti-Interleukin-3 IL3 Antibody catalog # PA1748. Tested in IHC, WB applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P08700

### **Technical Details**

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human IL-3.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized



888-466-3604 | support@bosterbio.com | www.bosterbio.com



Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat Western blot, 0.1-0.5ug/ml, Human




## Anti-Interleukin-3 IL3 Antibody (PA1748) Images



Anti-IL-3 antibody, PA1748, Western blotting All lanes: Anti IL-3 (PA1748) at 0.5ug/ml

Lane 1: Recombinant Human IL-3 Protein 10ng Lane 2: Recombinant Human IL-3 Protein 5ng Lane 3: Recombinant Human IL-3 Protein 2.5ng

Predicted bind size: 15KD Observed bind size: 15KD



Anti-IL-3 antibody, PA1748, IHC(P) IHC(P): Human Tonsil Tissue

# Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Interleukin-3 IL3 Antibody