

Anti-Transglutaminase 2/TGM2 Antibody

Catalog Number: PA2153

About TGM2

Tissue transglutaminase, also called TGC, is a 78-kDa, calcium dependent enzyme of the protein-glutamine gamma-glutamyltransferases family. By fluorescence in situ hybridization (FISH), this gene is mapped in 20q11.23. Transglutaminases are enzymes that catalyze the crosslinking of proteins by epsilon-gamma glutamyl lysine isopeptide bonds. While the primary structure of transglutaminases is not conserved, they all have the same amino acid sequence at their active sites and their activity is calcium-dependent. The protein encoded by this gene acts as a monomer, is induced by retinoic acid, and appears to be involved in apoptosis. Finally, the encoded protein is the autoantigen implicated in celiac disease.

Overview


Product Name	Anti-Transglutaminase 2/TGM2 Antibody
Reactive Species	Human, Rat
Description	Boster Bio Anti-Transglutaminase 2/TGM2 Antibody catalog # PA2153. Tested in IHC, ICC, WB applications. This antibody reacts with Human, Rat.
Application	IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Thimerosal, 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P21980

Technical Details


Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human TGM2, different from the related mouse sequence by three amino acids, and from the related rat sequence by four amino acids.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat</p> <p>Immunocytochemistry , 0.5-1ug/ml, Human, -</p> <p>Western blot, 0.1-0.5ug/ml, Human, Rat</p>

Anti-Transglutaminase 2/TGM2 Antibody (PA2153) Images



Anti-TGM2 antibody, PA2153, Western blotting
Lane 1: JURKAT Cell Lysate
Lane 2: HELA Cell Lysate



Anti-TGM2 antibody, PA2153, IHC(P)
IHC(P): Human Lung Cancer Tissue

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Transglutaminase 2/TGM2 Antibody