

Anti-Glutamate receptor 3/GRIA3 Antibody

Catalog Number: PA2249

About GRIA3

Glutamate receptor 3 is a protein that in humans is encoded by the GRIA3 gene. This gene belongs to a family of alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate (AMPA) receptors. It is mapped to Xq25. Glutamate receptors are the predominant excitatory neurotransmitter receptors in the mammalian brain and are activated in a variety of normal neurophysiologic processes. These receptors are heteromeric protein complexes with multiple subunits, each possessing transmembrane regions, and all arranged to form a ligand-gated ion channel. The classification of glutamate receptors is based on their activation by different pharmacologic agonists. In addition to that, GRIA3 has been shown to interact with GRIP1 and PICK1.

Overview

Product Name	Anti-Glutamate receptor 3/GRIA3 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Glutamate receptor 3/GRIA3 Antibody catalog # PA2249. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg Thimerosal, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P42263

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human GRIA3, identical to the related mouse and rat sequences.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized






BOSTER
antibody and ELISA experts

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Mouse, Rat, Human



Anti-Glutamate receptor 3/GRIA3 Antibody (PA2249) Images



Anti-GRIA3 antibody, PA2249, All Western blotting

All lanes: Anti-GRIA3 (PA2249) at 0.5ug/ml Lane 1: Rat Brain Tissue Lysate at 40ug Lane 2: Mouse Brain Tissue Lysate at 40ug

Predicted bind size: 101KD Observed bind size: 101KD

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Glutamate receptor 3/GRIA3 Antibody