

## Anti-Apg7/ATG7 Antibody

Catalog Number: PA2261

### About ATG7

Autophagy-related protein 7 is a protein that in humans is encoded by the ATG7 gene. It is mapped to 3p25.3. This gene was identified based on homology to *Pichia pastoris* GSA7 and *Saccharomyces cerevisiae* APG7. In the yeast, the protein appears to be required for fusion of peroxisomal and vacuolar membranes. The protein also shows homology to the ATP-binding and catalytic sites of the E1 ubiquitin activating enzymes. ATG7 is essential for the Apg12 conjugation system that mediates membrane fusion in autophagy. It is found that when nutrients are limited, ATG7 can regulate p53-dependent cell cycle and cell death pathways.

### Overview

Product Name	Anti-Apg7/ATG7 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Apg7/ATG7 Antibody catalog # PA2261. Tested in Flow Cytometry, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	O95352

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Apg7, different from the related mouse sequence by one amino acid, and from the related rat sequence by two amino acids.
Predicted Reactive Species	Chicken
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p> <p>Flow Cytometry, 1-3 ug/1x10<sup>6</sup> cells, Human</p>

## Anti-Apg7/ATG7 Antibody (PA2261) Images




Figure 1. Western blot analysis of Apg7/ATG7 using anti-Apg7/ATG7 antibody (PA2261). Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,  
Lane 2: human HepG2 whole cell lysates,  
Lane 3: human K562 whole cell lysates,  
Lane 4: human Jurkat whole cell lysates,  
Lane 5: rat kidney tissue lysates,  
Lane 6: mouse kidney tissue lysates,  
Lane 7: mouse spleen tissue lysates,  
Lane 8: mouse SP2/0 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-Apg7/ATG7 antigen affinity purified polyclonal antibody (Catalog # PA2261) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for Apg7/ATG7 at approximately 78 kDa. The expected band size for Apg7/ATG7 is at 78 kDa.




Figure 2. Flow Cytometry analysis of Hela cells using anti-Apg7/ATG7 antibody (PA2261).

Overlay histogram showing Hela cells stained with PA2261 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-Apg7/ATG7 Antibody (PA2261, 1 ug/1x10<sup>6</sup> cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10 ug/1x10<sup>6</sup> cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1 ug/1x10<sup>6</sup>) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

## 2 Publications Citing This Product

1. PubMed ID: 29552291, Erbin exerts a protective effect against inflammatory bowel disease by suppressing autophagic cell death
2. PubMed ID: 27383629, Nrf2 signalling and autophagy are involved in diabetes mellitus-induced defects in the development of mouse placenta

Visit [bosterbio.com/anti-apg7-antibody-pa2261-boster.html](http://bosterbio.com/anti-apg7-antibody-pa2261-boster.html) to see all 2 publications.

**Submit a product review to Biocompare.com**

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



## Anti-Apg7/ATG7 Antibody