

## Anti-Transcription factor SOX-2 SOX2 Antibody

Catalog Number: PA2284

### About SOX2

SRY (sex determining region Y)-box 2, also known as SOX2, is a transcription factor that is essential for maintaining self-renewal, or pluripotency of undifferentiated embryonic stem cells. Sox2 is a member of the Sox family of transcription factors, which have been shown to play key roles in many stages of mammalian development. This gene is mapped to 3q26.33. It is found that SOX2 can regulate OCT3/4 expression and maintains ES pluripotency through upstream transcription factors. SOX2 is identified as a lineage-survival oncogene in lung and esophageal squamous cell carcinoma. In addition to those, SOX2 has a critical role in maintenance of embryonic and neural stem cells and holds great promise in research involving induced pluripotency, an emerging and very promising field of regenerative medicine.

### Overview

Product Name	Anti-Transcription factor SOX-2 SOX2 Antibody
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Transcription factor SOX-2 SOX2 Antibody catalog # PA2284. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 4 mg Trehalose, 0.9 mg NaCl and 0.2 mg Na <sub>2</sub> HPO <sub>4</sub> .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P48431

### Technical Details

Immunogen	A synthetic peptide corresponding to a sequence in the middle region of human SOX2, different from the related mouse and rat sequence by one amino acid.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Immunohistochemistry (Paraffin-embedded Section), 2-5ug/ml, Human, Mouse, Rat, By Heat</p> <p>Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p>

## Anti-Transcription factor SOX-2 SOX2 Antibody (PA2284) Images




Figure 1. Western blot analysis of SOX2 using anti-SOX2 antibody (PA2284).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human U251 whole cell lysates,

Lane 2: human 293T whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SOX2 antigen affinity purified polyclonal antibody (Catalog # PA2284) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SOX2 at approximately 36 kDa. The expected band size for SOX2 is at 34 kDa.




Figure 2. IHC analysis of SOX2 using anti-SOX2 antibody (PA2284).

SOX2 was detected in a paraffin-embedded section of human laryngeal squamous cell carcinoma tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOX2 Antibody (PA2284) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.





Figure 3. IHC analysis of SOX2 using anti-SOX2 antibody (PA2284).

SOX2 was detected in a paraffin-embedded section of mouse brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOX2 Antibody (PA2284) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

Figure 4. IHC analysis of SOX2 using anti-SOX2 antibody (PA2284).



SOX2 was detected in a paraffin-embedded section of rat brain tissue. Heat mediated antigen retrieval was performed in EDTA buffer (pH 8.0, epitope retrieval solution). The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 2 ug/ml rabbit anti-SOX2 Antibody (PA2284) overnight at 4°C. Peroxidase Conjugated Goat Anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using HRP Conjugated Rabbit IgG Super Vision Assay Kit (Catalog # SV0002) with DAB as the chromogen.

## 2 Publications Citing This Product

1. PubMed ID: 25888093, Cheng Z, Wang Hz, Li X, Wu Z, Han Y, Li Y, Chen G, Xie X, Huang Y, Du Z, Zhou Y. J Exp Clin Cancer Res. 2015 Mar 26;34:27. Doi: 10.1186/S13046-015-0142-9. Microrna-184 Inhibits Cell Proliferation And Invasion, And Specifically Targets Tnfaip2 In G...
2. PubMed ID: 29531534, Yi H, Xie B, Liu B, Wang X, Xu L, Liu J, Li M, Zhong X, Peng F. Stem Cells Int. 2018 Jan 24;2018:3628578. doi: 10.1155/2018/3628578. eCollection 2018. Derivation and Identification of Motor Neurons from Human Urine-Derived Induced Pluripotent Stem...

Visit [bosterbio.com/anti-sox2-antibody-pa2284-boster.html](http://bosterbio.com/anti-sox2-antibody-pa2284-boster.html) to see all 2 publications.

## Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Transcription factor SOX-2 SOX2 Antibody