

Anti-E3 SUMO-protein ligase PIAS4 Antibody Picoband™

Catalog Number: PB10082

About PIAS4

E3 SUMO-protein ligase PIAS4, also known as protein inhibitor of activated STAT protein 4 (PIAS4) or protein inhibitor of activated STAT protein gamma (PIASg or PIASy), is an enzyme that in humans is encoded by the PIAS4 gene. This gene is mapped to 19p13.3. This gene plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway, the p53/TP53 pathway, the Wnt pathway and the steroid hormone signaling pathway. It also functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. This gene involved in gene silencing.

Overview

Product Name	Anti-E3 SUMO-protein ligase PIAS4 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-E3 SUMO-protein ligase PIAS4 Antibody Picoband™ catalog # PB10082. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q8N2W9

Technical Details

Immunogen	A synthetic peptide corresponding to a sequence at the N-terminus of human PIAS4, different from the related mouse sequence by two amino acids.
Predicted Reactive Species	Bovine, Canine, Horse, Monkey
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p>

Anti-E3 SUMO-protein ligase PIAS4 Antibody Picoband™ (PB10082) Images




Figure 1. Western blot analysis of PIAS4 using anti-PIAS4 antibody (PB10082).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat testis tissue lysates,
Lane 2: mouse testis tissue lysates,
Lane 3: HELA whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PIAS4 antigen affinity purified polyclonal antibody (Catalog # PB10082) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PIAS4 at approximately 57 kDa. The expected band size for PIAS4 is at 57 kDa.

Submit a product review to [Biocompare.com](https://www.biocompare.com)

Submit a review of this product to [Biocompare.com](https://www.biocompare.com) to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-E3 SUMO-protein ligase PIAS4 Antibody™