

Anti-Csk Antibody Picoband™

Catalog Number: PB9102

About CSK

CSK also known as C-Src kinase, is a non-receptor protein tyrosine kinase protein that in humans is encoded by the CSK gene. It is mapped to 15q24.1. CSK suppresses activity of the Src family of protein kinases by phosphorylation of Src family members at a conserved C-terminal site in Src. Csk's control of the Src family activity is widely thought to be central to regulation of the immune response. CSK can downregulate tyrosine kinase activity of the SRC oncoprotein through tyrosine phosphorylation of the SRC carboxy terminus. Since cell transformation by SRC oncoproteins is caused by various mechanisms that interfere with this phosphorylation, the CSK gene might function as an antioncogene.

Overview

Product Name	Anti-Csk Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Csk Antibody Picoband™ catalog # PB9102. Tested in IHC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg Na ₃ N.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P41240

Technical Details

Immunogen	E.coli-derived human CSK recombinant protein (Position: S2-G204). Human CSK shares 99% amino acid (aa) sequence identity with both mouse and rat CSK.
Predicted Reactive Species	Bovine, Horse, Monkey, Rabbit
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat</p>

Anti-Csk Antibody Picoband™ (PB9102) Images



Figure 1. Western blot analysis of CSK using anti-CSK antibody (PB9102).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. Lane 1: recombinant human CSK protein 0.5 ng. After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CSK antigen affinity purified polyclonal antibody (Catalog # PB9102) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CSK at approximately 47 kDa. The expected band size for CSK is at 47 kDa.





Figure 2. Western blot analysis of CSK using anti-CSK antibody (PB9102).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample under of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: rat testis tissue lysates,
Lane 2: rat thymus tissue lysates,
Lane 3: mouse liver tissue lysates,
Lane 4: human Hela whole cell lysates,
Lane 5: human Jurkat whole cell lysates,
Lane 6: human A549 whole cell lysates,
Lane 7: human MCF-7 whole cell lysates,
Lane 8: mouse NIH/3T3 whole cell lysates,
Lane 9: mouse NEURO whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CSK antigen affinity purified polyclonal antibody (Catalog # PB9102) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CSK at approximately 51 kDa. The expected band size for CSK is at 51 kDa.

Figure 3. IHC analysis of CSK using anti-CSK antibody (PB9102). CSK was detected in paraffin-embedded section of human tonsil tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat



serum. The tissue section was then incubated with 1ug/ml rabbit anti-CSK Antibody (PB9102) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.




Figure 4. IHC analysis of Csk using anti-Csk antibody (PB9102).

Csk was detected in paraffin-embedded section of human tonsil tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-Csk Antibody (PB9102) overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Streptavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

1 Publications Citing This Product

1. PubMed ID: , Multiple Signaling Roles of CD3epsilon and Its Application in CAR-T Cell Therapy

Visit bosterbio.com/anti-csk-picoband-trade-antibody-pb9102-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-Csk Antibody TM