

Anti-RIP/RIPK1 Antibody Picoband™

Catalog Number: PB9116

About RIPK1

RIPK1, also known as RIP or RIP1, is an enzyme that in humans is encoded by the RIPK1 gene. It is mapped to 6p25.2. RIPK1 is a key signaling molecule in the programmed necrosis pathway, which plays important roles in development, tissue damage response, and antiviral immunity. RIPK1 is known to have function in a variety of cellular pathways including the NF-kappaB pathway and programmed necrotic cell death (necroptosis). The kinase domain, while important for necroptotic (programmed necrotic) functions, it appears dispensable for other lethal, as well as pro-survival roles. Also, proteolytic processing of RIPk1, through both caspase-dependent and -independent mechanisms, triggers lethality that is dependent on the generation of one or more specific C-terminal cleavage product (s) of RIPk1 upon stress.

Overview

Product Name	Anti-RIP/RIPK1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-RIP/RIPK1 Antibody Picoband™ catalog # PB9116. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q13546

Technical Details

Immunogen	E.coli-derived human RIP recombinant protein (Position: K316-N671). Human RIP shares 65% amino acid (aa) sequence identity with mouse RIP.
Predicted Reactive Species	Bovine, Horse, Monkey, Rabbit
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human

Anti-RIP/RIPK1 Antibody Picoband™ (PB9116) Images



Figure 1. Western blot analysis of RIP using anti-RIP antibody (PB9116).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours.

Lane 1: recombinant human RIP protein 0.5 ng.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RIP antigen affinity purified polyclonal antibody (Catalog # PB9116) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for RIP at approximately 38 kDa. The expected band size for RIP is at 38 kDa.




Figure 2. Western blot analysis of RIP using anti-RIP antibody (PB9116).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Jurkat whole cell lysates,

Lane 2: human 22RV1 whole cell lysates,

Lane 3: human MCF-7 whole cell lysates,

Lane 4: human Hela whole cell lysates,

Lane 5: human A549 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes. Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-RIP antigen affinity purified polyclonal antibody (Catalog # PB9116) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for RIP at approximately 76 kDa. The expected band size for RIP is at 76 kDa.

3 Publications Citing This Product

1. PubMed ID: -, Smad3-Targeted Therapy Protects against Cisplatin-Induced AKI by Attenuating Programmed Cell Death and Inflammation via a NOX4-Dependent Mechanism. Qin Yang, Li Gao, Xiao-wei Hu, Jia-nan Wang, Yao Zhang, Yu-hang Dong, Hui Yao Lan, Xiao-ming Meng

2. PubMed ID: 27464624, Bifidobacterial recombinant thymidine kinase-ganciclovir gene therapy system induces FasL and TNFR2 mediated antitumor apoptosis in solid tumors

3. PubMed ID: 25674205, Chen Yf, Zhao Zq, Wu Zm, Zou Zy, Luo Xj, Li J, Xie C, Liang Y. Int J Clin Exp Pathol. 2014 Dec 1;7(12):8411-20. Ecollection 2014. The Role Of Rip1 And Rip3 In The Development Of Aplastic Anemia Induced By Cyclophosphamide And Busulphan In Mice.

Visit bosterbio.com/anti-rip-picoband-trade-antibody-pb9116-boster.html to see all 3 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-RIP/RIPK1 Antibody [™]