

Anti-SIRT1 Antibody Picoband™

Catalog Number: PB9159

About SIRT1

Sirtuin 1, also known as SIR2L1 or SIRT1, is a protein that in humans is encoded by the SIRT1 gene. It is mapped to 10q21.3. Sirtuin 1 is a member of the sirtuin family of proteins, homologs of the Sir2 gene in *S. cerevisiae*. Members of the sirtuin family are characterized by a sirtuin core domain and grouped into four classes. Sirtuin 1 is downregulated in cells that have high insulin resistance and inducing its expression increases insulin sensitivity, suggesting the molecule is associated with improving insulin sensitivity. Furthermore, Sirtuin 1 was shown to de-acetylate and affect the activity of both members of the PGC1-alpha/ERR-alpha complex, which are essential metabolic regulatory transcription factors.

Overview

Product Name	Anti-SIRT1 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-SIRT1 Antibody Picoband™ catalog # PB9159. Tested in IF, ICC, WB applications. This antibody reacts with Human.
Application	IF, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q96EB6

Technical Details

Immunogen	E.coli-derived human SIRT1 recombinant protein (Position: R112-K311). Human SIRT1 shares 90% amino acid (aa) sequence identity with mouse SIRT1.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human Immunocytochemistry/Immunofluorescence, 5 ug/ml, Human

Anti-SIRT1 Antibody Picoband™ (PB9159) Images




Figure 1. Western blot analysis of SIRT1 using anti-SIRT1 antibody (PB9159).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human HepG2 whole cell lysates,

Lane 2: human MCF-7 whole cell lysates,

Lane 3: human SiHa whole cell lysates,

Lane 4: human Daudi whole cell lysates,

Lane 5: human K562 whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5

hour at RT. The membrane was incubated with rabbit anti-

SIRT1 antigen affinity purified polyclonal antibody (Catalog

PB9159) at 0.5 ug/mL overnight at 4°C, then washed with

TBS-0.1%Tween 3 times with 5 minutes each and probed

with a goat anti-rabbit IgG-HRP secondary antibody at a

dilution of 1:5000 for 1.5 hour at RT. The signal is developed

using an Enhanced Chemiluminescent detection (ECL) kit

(Catalog # EK1002) with Tanon 5200 system. A specific

band was detected for SIRT1 at approximately 120 kDa. The

expected band size for SIRT1 is at 82 kDa.




Figure 2. IF analysis of SIRT1 using anti-SIRT1 antibody (PB9159).

SIRT1 was detected in an immunocytochemical section of A431 cells. Enzyme antigen retrieval was performed using IHC enzyme antigen retrieval reagent (AR0022) for 15 mins.

The cells were blocked with 10% goat serum. And then

incubated with 5 ug/mL rabbit anti-SIRT1 Antibody (PB9159)

overnight at 4°C. DyLight®488 Conjugated Goat Anti-Rabbit

IgG (BA1127) was used as secondary antibody at 1:100

dilution and incubated for 30 minutes at 37°C. The section

was counterstained with DAPI. Visualize using a fluorescence

microscope and filter sets appropriate for the label used.

1 Publications Citing This Product

1. PubMed ID: 25162824, Hou S, Zheng F, Li Y, Gao L, Zhang J. Int J Mol Sci. 2014 Aug 26;15(9):15026-43. Doi: 10.3390/ijms150915026. The Protective Effect Of Glycyrrhizic Acid On Renal Tubular Epithelial Cell Injury Induced By High Glucose.

Visit bosterbio.com/anti-sirt1-picoband-trade-antibody-pb9159-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-SIRT1 Antibody [™]