

Anti-Fas Antibody Picoband™

Catalog Number: PB9252

About FAS

FAS (also known as surface antigen APO1 or CD95) is a member of the tumour-necrosis receptor factor family of death receptors. It acts as an inducer of both neurite growth in vitro and accelerated recovery after nerve injury in vivo. FAS antigen is expressed and functional on papillary thyroid cancer cells and this may have potential therapeutic significance. The FAS antigen shows structural homology with a number of cell surface receptors, including tumor necrosis factor (TNF) receptors and the low-affinity nerve growth factor receptor (NGFR) and it is mapped to 10q24.1. The FAS and FASL system plays a key role in regulating apoptotic cell death and corruption of this signalling pathway has been shown to participate in immune escape and tumorigenesis.

Overview

Product Name	Anti-Fas Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-Fas Antibody Picoband™ catalog # PB9252. Tested in IHC, WB applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P25445

Technical Details

Immunogen	E.coli-derived human Fas recombinant protein (Position: Q26-N173). Human Fas shares 55% and 59% amino acid (aa) sequences identity with mouse and rat Fas, respectively.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized






888-466-3604 | support@bosterbio.com | www.bosterbio.com

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat Western blot, 0.1-0.5ug/ml, Human




Anti-Fas Antibody Picoband™ (PB9252) Images



Anti-FAS Picoband antibody, PB9252, Western blotting

All lanes: Anti FAS (PB9252) at 0.5ug/ml WB: Recombinant Human FAS Protein 0.5ng


Predicted bind size: 35KD Observed bind size: 35KD



Anti-FAS Picoband antibody, PB9252, Western blotting

All lanes: Anti FAS (PB9252) at 0.5ug/ml Lane 1: HELA Whole Cell Lysate at 40ug Lane 2: JURKAT Whole Cell Lysate at 40ug Lane 3: A549 Whole Cell Lysate at 40ug Lane 4: SMMC Whole Cell Lysate at 40ug Lane 5: K562 Whole Cell Lysate at 40ug

Predicted bind size: 37KD Observed bind size: 37KD



Anti-FAS Picoband antibody, PB9252, IHC(P) IHC(P): Human Lung Cancer Tissue

24 Publications Citing This Product

- 1. PubMed ID: PMID:26617948, Significance of Fas and FasL protein expression in cardiac carcinoma and local lymph node tissues
- 2. PubMed ID: 10.3748/wjg.v16.i10.1274, Clinical significance of Fas and FasL protein expression in gastric carcinoma and local lymph node tissues
- 3. PubMed ID: 10.1039/C6F000071A, The effect of geniste on Abeta25-35-induced PC12 cell apoptosis through the JNK-dependent Fas pathway

Visit bosterbio.com/anti-fas-picoband-trade-antibody-pb9252-boster.html to see all 24 publications.

Submit a product review to Biocompare.com











Anti-Fas Antibody ™