

Anti-CaV1.3/CACNA1D Antibody Picoband™

Catalog Number: PB9329

About CACNA1D

CACNA1D is also known as PASNA, SANDD or Cav1.3. Voltage-dependent calcium channels mediate the entry of calcium ions into excitable cells, and are also involved in a variety of calcium-dependent processes, including muscle contraction, hormone or neurotransmitter release, and gene expression. Calcium channels are multisubunit complexes composed of alpha-1, beta, alpha-2/delta, and gamma subunits. The channel activity is directed by the pore-forming alpha-1 subunit, whereas the others act as auxiliary subunits regulating this activity. The distinctive properties of the calcium channel types are related primarily to the expression of a variety of alpha-1 isoforms, namely alpha-1A, B, C, D, E, and S. This gene encodes the alpha-1D subunit. Several transcript variants encoding different isoforms have been found for this gene.

Overview

Product Name	Anti-CaV1.3/CACNA1D Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-CaV1.3/CACNA1D Antibody Picoband™ catalog # PB9329. Tested in Flow Cytometry, IHC, ICC, WB applications. This antibody reacts with Human, Mouse, Rat.
Application	Flow Cytometry, IHC, ICC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q01668

Technical Details

Immunogen	E.coli-derived human CaV1.3 recombinant protein (Position: M1-I180). Human CaV1.3 shares 99% and 98% amino acid (aa) sequence identity with mouse and rat CaV1.3, respectively.
Predicted Reactive Species	Bovine
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(F) and ICC.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized

Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.
Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Mouse, Rat, Human Immunohistochemistry (Frozen Section), 0.5-1ug/ml, Human Immunocytochemistry, 0.5-1ug/ml, Human Flow Cytometry, 1-3ug/1x10⁶ cells, Human

Anti-CaV1.3/CACNA1D Antibody Picoband™ (PB9329) Images



Figure 1. Western blot analysis of CAV1.3 using anti-CAV1.3 antibody (PB9329).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 50ug of sample under reducing conditions.

Lane 1: Rat Brain Tissue Lysate,

Lane 2: Mouse Brain Tissue Lysate.

After Electrophoresis, proteins were transferred to a Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-CAV1.3 antigen affinity purified polyclonal antibody (Catalog # PB9329) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for CAV1.3 at approximately 245KD. The expected band size for CAV1.3 is at 245KD.




Figure 2. Flow Cytometry analysis of U-87 cells using anti-CaV1.3 antibody (PB9329).

Overlay histogram showing U-87 cells stained with PB9329 (Blue line). The cells were blocked with 10% normal goat serum. And then incubated with rabbit anti-CaV1.3 Antibody (PB9329, 1ug/1x10⁶ cells) for 30 min at 20°C. DyLight®488 conjugated goat anti-rabbit IgG (BA1127, 5-10ug/1x10⁶ cells) was used as secondary antibody for 30 minutes at 20°C. Isotype control antibody (Green line) was rabbit IgG (1ug/1x10⁶) used under the same conditions. Unlabelled sample (Red line) was also used as a control.

1 Publications Citing This Product

1. PubMed ID: 28656274, Guo, Y., Cui, L., Jiang, S., Zhang, A., & Jiang, S. (2017). Proteomics of acute heart failure in a rat post-myocardial infarction model. Molecular Medicine Reports, 16(2), 1946-1956. Advance online publication. doi: 10.3892/mmr.2017.6820

Visit bosterbio.com/anti-cav1-3-picoband-trade-antibody-pb9329-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

