

Anti-TPA Tissue Plasminogen Activator/PLAT Antibody Picoband™

Catalog Number: PB9345

About PLAT

PLAT is also known as tPA. This gene encodes tissue-type plasminogen activator, a secreted serine protease which converts the proenzyme plasminogen to plasmin, a fibrinolytic enzyme. Tissue-type plasminogen activator is synthesized as a single chain which is cleaved by plasmin to a two chain disulfide linked protein. This enzyme plays a role in cell migration and tissue remodeling. Increased enzymatic activity causes hyperfibrinolysis, which manifests as excessive bleeding; decreased activity leads to hypofibrinolysis which can result in thrombosis or embolism. Alternative splicing of this gene results in multiple transcript variants encoding different isoforms.

Overview

Product Name	Anti-TPA Tissue Plasminogen Activator/PLAT Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-TPA Tissue Plasminogen Activator/PLAT Antibody Picoband™ catalog # PB9345. Tested in WB applications. This antibody reacts with Human.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na ₂ HPO ₄ , 0.05mg NaN ₃ .
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P00750

Technical Details

Immunogen	E.coli-derived human TPA recombinant protein (Position: H366-P562). Human TPA shares 83% and 84% amino acid (aa) sequence identity with mouse and rat TPA, respectively.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.

Purification	Immunogen affinity purified.
Suggested Dilutions	<p>Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.</p> <p>If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.</p> <p>Some PubMed article(s) citing the expression level of this target are as follows:</p> <p>Boster Bio's internal QC testing used:</p> <p>Western blot, 0.1-0.5ug/ml, Human</p>

Anti-TPA Tissue Plasminogen Activator/PLAT Antibody Picoband™ (PB9345) Images




Figure 1. Western blot analysis of PLAT using anti-PLAT antibody (PB9345).

Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each lane was loaded with 30 ug of sample under reducing conditions.

Lane 1: human Hela whole cell lysates,

Lane 2: human PC-3 whole cell lysates,

Lane 3: human U87 whole cell lysates,

Lane 4: human SH-SY5Y whole cell lysates.

After electrophoresis, proteins were transferred to a nitrocellulose membrane at 150 mA for 50-90 minutes.

Blocked the membrane with 5% non-fat milk/TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PLAT antigen affinity purified polyclonal antibody (Catalog # PB9345) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:5000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PLAT at approximately 63-69 kDa. The expected band size for PLAT is at 63 kDa.

1 Publications Citing This Product

1. PubMed ID: 34082810, Wang J,Li Y,Pan L,Li J,Yu Y,Liu B,Zubair M,Wei Y,Pillay B,Olaniran AO,Chiliza TE,Shao G,Feng Z,Xiong Q.Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) moonlights as an adhesin in Mycoplasma hyorhinis adhesion to epithelial cells as well as a plasminogen receptor mediating extracellular matrix degradation.Vet Res.2021 Jun 3;52(1):80.doi:10.1186/s13567-021-00952-8. PMID:34082810;PMCID:PMC8173509.

Visit bosterbio.com/anti-tpa-picoband-trade-antibody-pb9345-boster.html to see all 1 publications.

Submit a product review to Biocompare.com

Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.



Anti-TPA Tissue Plasminogen Activator/PLAT Antibody™