

Anti-SerpinB2 Antibody Picoband™

Catalog Number: PB9355

About SERPINB2

SERPINB2 is also known as PAI or PAI2. Plasminogen activator inhibitor-2 (placental PAI) is acoagulation factor that inactivates tPA and urokinase. It is present in most cells, especially monocytes/macrophages. PAI-2 exists in two forms, a 60-kDa extracellular glycosylated form and a 43-kDa intracellular form. It is present only at detectable quantities in blood during pregnancy, as it is produced by the placenta, and may explain partially the increased rate of thrombosisduring pregnancy. The majority of expressed PAI-2 remains unsecreted due to the presence of an inefficient internal signal peptide.

Overview

Product Name	Anti-SerpinB2 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-SerpinB2 Antibody Picoband™ catalog # PB9355. Tested in IHC, WB applications. This antibody reacts with Human.
Application	IHC, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P05120

Technical Details

Immunogen	E.coli-derived human SerpinB2 recombinant protein (Position: M1-K180). Human SerpinB2 shares 70% and 66% amino acid (aa) sequence identity with mouse and rat SerpinB2, respectively.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Immunohistochemistry (Paraffin-embedded Section), 0.5-1ug/ml, Human, By Heat Western blot, 0.1-0.5ug/ml, Human




Anti-SerpinB2 Antibody Picoband™ (PB9355) Images

130KD -100KD -70KD -55KD -35KD -25KD -

Anti-SerpinB2 Picoband antibody, PB9355, Western blotting

All lanes: Anti SerpinB2 (PB9355) at 0.5ug/ml WB: Human Placenta Tissue Lysate at 50ug

Predicted bind size: 47KD Observed bind size: 47KD



Anti-SerpinB2 Picoband antibody, PB9355,IHC(P) IHC(P): Human Placenta Tissue

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-SerpinB2 Antibody ™