

Anti-SFRP2 Antibody Picoband™

Catalog Number: PB9362

About SFRP2

SFRP2 is also known as FRP-2, SARP1 or SDF-5. It is Expressed in adipose tissue, heart, brain, skeletal muscle, pancreas, thymus, prostate, testis, ovary, small intestine and colon. Highest levels in adipose tissue, small intestine and colon. This gene encodes a member of the SFRP family that contains a cysteine-rich domain homologous to the putative Wnt-binding site of Frizzled proteins. SFRPs act as soluble modulators of Wnt signaling. Methylation of this gene is a potential marker for the presence of colorectal cancer.

Overview

Product Name	Anti-SFRP2 Antibody Picoband™
Reactive Species	Human
Description	Boster Bio Anti-SFRP2 Antibody Picoband™ catalog # PB9362. Tested in ELISA, WB applications. This antibody reacts with Human.
Application	ELISA, WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	Q96HF1

Technical Details

Immunogen	E.coli-derived human SFRP2 recombinant protein (Position: L104-C295). Human SFRP2 shares 99% amino acid (aa) sequence identity with mouse SFRP2.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.



BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit. If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples. Some PubMed article(s) citing the expression level of this target are as follows: Boster Bio's internal QC testing used: Western blot, 0.1-0.5ug/ml, Human, - ELISA, 0.1-0.5ug/ml, Human



Anti-SFRP2 Antibody Picoband™ (PB9362) Images



Anti-SFRP2 Picoband antibody, PB9362, Western blotting

All lanes: Anti SFRP2 (PB9362) at 0.5ug/ml Lane 1: COLO320 Whole Cell Lysate at 40ug Lane 2: SW620 Whole Cell Lysate at 40ug

Predicted bind size: 33KD Observed bind size: 33KD

Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-SFRP2 Antibody ™