

# Anti-Caspase-2/CASP2 Antibody Picoband™

Catalog Number: PB9368

#### **About CASP2**

CASP2 is equal to Caspase-2. And Caspase-2, which is involved in stress-induced apoptosis, is recruited into a large protein complex, the molecular composition of which remains elusive. It is showed that activation of caspase-2 occurs in a complex that contains the death domain-containing protein PIDD, whose expression is induced by p53, and the adaptor protein RAIDD. Increased PIDD expression resulted in spontaneous activation of caspase-2 and sensitization to apoptosis by genotoxic stimuli. Caspase-2 acts both as a positive and negative cell death effector, depending upon cell lineage and stage of development.

#### Overview

Product Name	Anti-Caspase-2/CASP2 Antibody Picoband™
Reactive Species	Human, Mouse, Rat
Description	Boster Bio Anti-Caspase-2/CASP2 Antibody Picoband™ catalog # PB9368. Tested in WB applications. This antibody reacts with Human, Mouse, Rat.
Application	WB
Clonality	Polyclonal
Formulation	Each vial contains 5mg BSA, 0.9mg NaCl, 0.2mg Na2HPO4, 0.05mg NaN3.
Storage Instructions	Store at -20°C for one year from date of receipt. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20°C for six months. Avoid repeated freeze-thaw cycles.
Host	Rabbit
Uniprot ID	P42575

#### **Technical Details**

Immunogen	A synthetic peptide corresponding to a sequence at the C-terminus of human Caspase-2, different from the related mouse and rat sequences by one amino acid.
Predicted Reactive Species	Hamster
Recommended Detection Systems	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (EK1002) for Western blot.
Cross Reactivity	No cross-reactivity with other proteins.
Isotype	Rabbit IgG
Form	Lyophilized
Concentration	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml.




# BOSTER BIOLOGICAL TECHNOLOGY 3942 B Valley Ave, Pleasanton, CA 94566

888-466-3604 | support@bosterbio.com | www.bosterbio.com

Purification	Immunogen affinity purified.
Suggested Dilutions	Dilute the sample so that the expected range of concentrations fall within the detection range of this kit.  If the expected range of concentration is unknown, a pilot test should be conducted to decide the optimal dilution ratio for your samples.  Some PubMed article(s) citing the expression level of this target are as follows:  Boster Bio's internal QC testing used:  Western blot, 0.1-0.5ug/ml, Human, Mouse, Rat



### Anti-Caspase-2/CASP2 Antibody Picoband™ (PB9368) Images



Anti-Caspase-2 Picoband antibody, PB9368, Western blotting

All lanes: Anti Caspase-2 (PB9368) at 0.5ug/ml Lane 1: Rat Lung Tissue Lysate at 50ug

Lane 2: Mouse Liver Tissue Lysate at 50ug Lane 3: A549 Whole Cell Lysate at 40ug Lane 4: PANC Whole Cell Lysate at 40ug

Lane 5: 293T Whole Cell Lysate at 40ug Predicted bind size: 18KD Observed bind size: 18KD

## Submit a product review to Biocompare.com





Submit a review of this product to Biocompare.com to receive a \$20 Amazon.com giftcard! Your reviews help your fellow scientists make the right decisions. Thank you for your contribution.

Anti-Caspase-2/CASP2 Antibody ™